



Flanders
State of
the Art

Annex: Monetisation of the MMG method

[update 2017]

WE MAKE
TOMORROW
BEAUTIFUL
OVAM

WWW.OVAM.BE

**Annex: Monetisation of
the MMG method (update
2017)**



Document description

1. *Title of publication*

Annex: Monetisation of the MMG method (update 2017)

2. *Responsible Publisher*

Danny Wille, OVAM, Stationsstraat 110, 2800 Mechelen

3. *Legal deposit number*

D/2018/5024/07

4. *Number of pages*

65

5. *Number of tables and figures*

35 tables and 1 figure

6. *Price**

7. *Date of publication*

January 2018

8. *Keywords*

monetisation, MMG determination method, indicators

9. *Summary*

This note describes the methods and data used to monetise the LCIA impacts as a further development of the approach used in MMG2012 (Debacker et al., 2012). First, we discuss some general changes to methods and data. Second, we discuss for each impact category the approach, data and results. We focus on new developments (i.e. MMG2014) compared to MMG2012, and refer to previous studies for the details. For some indicators updates have been made anno 2017.

10. *Steering committee and/or author*

Leo De Nocker, VITO - Wim Debacker, VITO

11. *Contact person(s)*

roos.servaes@ovam.be, Roos Servaes
denocker@vito.be, Leo De Nocker – wim.debacker@vito.be, Wim Debacker

12. *Other publications about the same subject*

Environmental profile of building elements
(http://www.ovam.be/sites/default/files/FILE1368696514672Environmental_profile_buildig_elements_LR.pdf)

Information from this document may be reproduced with due acknowledgement of the source.

Most OVAM publications can be consulted and/or downloaded on OVAM's website: <http://www.ovam.be>

* Price changes reserved.

Table of contents

	Document description	3
1	Introduction	7
2	Summary overview of methods and results	9
2.1	Methods	9
2.2	Results	9
3	Methodology: selection of data and uncertainties	13
3.1	Selection of methods and data	13
3.1.1	Indicators for social costs based on damage costs	13
3.1.2	Indicators for social costs based on prevention costs	14
3.1.3	Selection of information for central estimate and low-high estimates	15
3.1.4	Comparison with literature	15
3.2	Discounting	15
3.3	Uncertainty	16
3.3.1	Background	16
3.3.2	Treatment of uncertainty in MMG2014	17
3.4	Comparison with the Dutch environment assessment methods for buildings	19
4	Monetary indicators for CEN indicators	21
4.1	Global Warming	21
4.1.1	Monetary value	21
4.1.2	Methods and data used	21
4.2	Depletion of the stratospheric ozone layer	24
4.2.1	Monetary value	24
4.2.2	Methods and data used	24
4.3	Acidification of land and water sources	24
4.3.1	Monetary value	24
4.3.2	Methods and data used	25
4.4	Eutrophication	25
4.4.1	Monetary value	25
4.4.2	Methods and data used	26
4.5	Formation of tropospheric ozone photochemical oxidants	27
4.5.1	Monetary value	27
4.5.2	Methods and data used	27
4.6	Abiotic depletion of non-fossil resources	28
4.6.1	Monetary value	28
4.6.2	Methods and data used	28
4.7	Abiotic depletion of fossil resources	31
4.7.1	Monetary value	31
4.7.2	Methods and data used	31
5	Monetary indicators for CEN+ indicators	33
5.1	Human toxicity, cancer effects	33
5.1.1	Monetary value	33
5.1.2	Methods and data used	33
5.1.3	Further explanation	33
5.1.4	Low and high values	35
5.2	Human toxicity, non-cancer effects	35
5.2.1	Monetary value	35
5.2.2	Methods and data used	35
5.3	Particulate matter	36
5.3.1	Monetary value	36
5.3.2	Methods and data used	36
5.3.3	Low and high values	37

5.4	Ionising radiation: human health effects	40
5.4.1	Monetary value	40
5.4.2	Methods and data used	40
5.5	Ecotoxicity, freshwater	41
5.5.1	Monetary value	41
5.5.2	Methods and data used	41
5.6	Water scarcity	43
5.6.1	Monetary value	43
5.6.2	Methods and data used	43
5.7	Land use occupation	46
5.7.1	Monetary value	46
5.7.2	Methods and data used	47
5.8	Land use transformation	50
5.8.1	Monetary value	50
5.8.2	Methods and data used	51
Annex 1:	List of tables	53
Annex 2:	List of figures	55
Annex 3:	Bibliography	57

1 Introduction

Objective: the analysis in Debacker et al. (2012) indicates that the monetisation step can have a big impact on the (aggregated) MMG results. Monetising impacts both reveals the uncertainty inherent in LCIA and adds uncertainty. In 2013-2014, we further developed the monetization step, especially to make the approach consistent with the indicators used in ILCD recommendations and CEN guidelines, and to clarify and narrow down the uncertainties.

This note describes the methods and data used to monetise the LCIA impacts as a further development of the approach used in MMG2012 (Debacker et al., 2012). First, we discuss some general changes to methods and data. Second, we discuss for each impact category the approach, data and results. We focus on new developments (i.e. MMG2014) compared to MMG2012, and refer to previous studies for the details. In the following paragraphs 'MMG2012' will refer to the original assessment method developed in 2012. With 'MMG2014' we will refer to new developments. In 2017, the impact category 'Ionising radiation: ecosystems' has been removed from the set of life cycle impact assessment methods and small updates have been done for the monetary values of the impact categories 'Global warming', 'Land use occupation', and 'Land use transformation'.

2 Summary overview of methods and results

2.1 Methods

The methods and data build further on methods and data selected in the context of the MMG2012 project (Debacker et al, 2012; De Nocker, 2012).

In the context of the current project, the methods and data are updated and adapted to account for:

- Consistency of the indicators for LCI inventory and LCIA impact assessment, in line with the CEN and CEN+ recommendations. The list of impacts to be covered and the selected metrics and LCIA models/methods are given in table 1 and 2 for the CEN and CEN+ indicators respectively. The LCIA models for the CEN indicators are based on the latest version of the EN 15804 standard (CEN 2013). The LCIA models for the CEN+ indicators are based on ILCD recommendations (JRC 2011), with the exception of the land use biodiversity indicators.
- For 'Land Transformation' and 'Land Occupation' the sub indicator 'biodiversity' has been added. Based on Allacker et al. (2014) and Debacker et al. (2012), land use impacts related to changes in biodiversity are significant for some processes and products within the built environment.
- In addition, the scope and characteristics of the LCIA models in these steps may have impacts for the selection of the monetary value.
- Update of new information related to monetization of environmental impacts.
- A more coherent approach to discounting, using a 3 % discount rate as the central estimate.
- A more consistent approach to deal with uncertainty for different impact categories and to present a central, low and high estimate.
- Monetary values of each environmental indicator have been determined for three regions: Western Europe, Flanders/Belgium and the rest of the world. Only the Western European values are taken into account for the publically available MMG method, because most processes related to the life cycle of building products are related to this geographical area. The monetary values for the Flemish/Belgian region and the rest of the world are determined for sensitivity analyses. The uncertainty related to the monetary value is the smallest for the Flemish/Belgian region (or equal to the Western European region and the rest of the world).

2.2 Results

Tables 1, 2, 3 and 4 give an overview of the results for Western Europe. In sections 4 and 5, we also report the values for Flanders/Belgium and for the 'rest of the world' category.

Environmental indicator (CEN)	Unit	Selected LCIA model
1. Global warming	kg CO2 eqv.	EN 15804+A1 *
2. Depletion of the stratospheric ozone layer	kg CFC-11 eqv.	EN 15804+A1 *
3. Acidification of land and water sources	kg SO2 eqv.	EN 15804+A1 *
4. Eutrophication	kg (PO4)3- eqv.	EN 15804+A1 *
5. Formation of tropospheric ozone photochemical oxidants	kg etheen eqv.	EN 15804+A1 *
6. Abiotic depletion of non-fossil resources	kg Sb** eqv.	EN 15804+A1 *
7. Abiotic depletion of fossil resources	MJ, net caloric value	EN 15804+A1 *

Table 1: Selected CEN environmental indicators, the corresponding units and LCIA models

* EN 15804+A1 = EN 15804+A1 (as used in CML version 2012)

** Sb: antimony

Environmental indicator (CEN+)	Unit	Selected LCIA model
8. Human toxicity		
a. cancer effects	CTUh ^{***}	Rosenbaum et al., 2008 (as used in USEtox)
b. non-cancer effects	CTUh ^{***}	Rosenbaum et al., 2008 (as used in USEtox)
9. Particulate matter	kg PM _{2,5} eqv.	Rabl et al, 2004 (as used in RiskPoll)
10. Ionising radiation, human health	kg U235 eqv.	Frishknecht et al., 2000 (as used in ReCiPe midpoint)
11. Ecotoxicity ¹ :		
a. terrestrial	-	-
b. freshwater	CTUe ^{****}	Rosenbaum et al., 2008 (as used in USEtox)
c. marine	-	-
12. Water scarcity	m ³ water eqv.	Frishknecht et al, 2006 (as used in Swiss Ecoscarcy – water)
13. Land use: occupation:		
a. soil organic matter	kg C deficit	Milà i Canals et al., 2007
b. biodiversity ²	m ² a	Köllner, 2000 + characterisation factors set on "1"
14. Land use: transformation		
a. soil organic matter	kg C deficit	Milà i Canals et al., 2007
b. biodiversity ³	m ²	Köllner, 2000 + characterisation factors set on "1" or "-1"

Table 2: Selected CEN+ environmental indicators, the corresponding units and LCIA models

Environmental indicator (CEN+) Unit Selected LCIA model

*** CTUh: comparative toxic units for humans

**** CTUe: comparative toxic units for ecosystems

¹ For terrestrial and marine ecotoxicity, the PEF Guide does not recommend an LCIA model

² Biodiversity impacts related to land use are not taken into account in the PEF. However, due to its importance in the built environment, the Köllner 2000 model as used in Eco-Indicator 99 (in PDF*m²yr and PDF*m²) is proposed as a best proxy to take into account biodiversity impacts related to land use for the individual MMG scoring. Impacts of land use occupation and land use transformation expressed in m²a and m² respectively have proven to be a better basis to calculate the related environmental costs. For this reason, the land use occupation and transformation processes (expressed per m²a and m² respectively) considered in the Köllner 2000 model are taken into account, but characterisation factors are set to "1" or "-1" for the calculation of the environmental costs and in this way used as an inventory method for biodiversity flows. For the calculation of the individual indicators the Köllner 2000 model as used in Eco-Indicator 99 (in PDF*m²yr and PDF*m²) is used

³ Biodiversity impacts related to land use are not taken into account in the PEF. However, due to its importance in the built environment, the Köllner 2000 model as used in Eco-Indicator 99 (in PDF*m²yr and PDF*m²) is proposed as a best proxy to take into account biodiversity impacts related to land use for the individual MMG scoring. Impacts of land use occupation and land use transformation expressed in m²a and m² respectively have proven to be a better basis to calculate the related environmental costs. For this reason, the land use occupation and transformation processes (expressed per m²a and m² respectively) considered in the Köllner 2000 model are taken into account, but characterisation factors are set to "1" or "-1" for the calculation of the environmental costs and in this way used as an inventory method for biodiversity flows. For the calculation of the individual indicators the Köllner 2000 model as used in Eco-Indicator 99 (in PDF*m²yr and PDF*m²) is used

Environmental indicator (CEN)	Unit	Central (€/unit)	Low (€/unit)	High (€/unit)
1. Global warming	kg CO2 eqv.	0.050	0.025	0.100
2. Depletion of the stratospheric ozone layer	kg CFC-11 eqv.	49.10	25	100
3. Acidification of land and water sources	kg SO2 eqv.	0.43	0.22	0.88
4. Eutrophication	kg (PO4) ³⁻ eqv.	20	6.60	60
5. Formation of tropospheric ozone photochemical oxidants	kg ethene eqv.	0.48	0	6.60
6. Abiotic depletion of non-fossil resources	kg Sb eqv.	1.56	0	6.23
7. Abiotic depletion of fossil resources	MJ, net caloric value	0	0	0.0065

Table 3: Overview of West-European monetary (central, low and high) values for the CEN indicators

Environmental indicator (CEN+)	Unit	Central (€/unit)	Low (€/unit)	High (€/unit)
8. Human toxicity				
a. cancer effects	CTUh	665109	166277	2660434
b. non-cancer effects	CTUh	144081	28816	720407
9. Particulate matter	kg PM2,5 eqv.	34	12.70	85
10. Ionising radiation, human health	kg U235 eqv.	9.7E-04	3.2E-04	2.9E-03
11. Ecotoxicity:				
a. terrestrial	-			
b. freshwater	CTUe	3.70E-05	7.39E-06	1.85E-04
c. marine	-			
12. Water scarcity	m ³ water eqv.	0.067	0.022	0.20
13. Land use: occupation:				
a. soil organic matter	kg C deficit	1.4E-06	3.4E-07	0.6E-05
b. biodiversity				
b1. urban: loss ES*	m ² .a	0.30	0.07	2.35
b2. agricultural	m ² .a	6.0E-03	1.5E-03	2.4E-02
b3. forest: biodiversity	m ² .a	2.2E-04	5.5E-05	8.8E-04
14. Land use: transformation				
a. soil organic matter	kg C deficit	1.4E-06	3.4E-07	0.6E-05
b. biodiversity				
b1. urban	m ²	n.a.	n.a.	n.a.
b2. agricultural	m ²	n.a.	n.a.	n.a.
b3. forest, excl. tropical	m ²	n.a.	n.a.	n.a.
b4. tropical rainforest	m ²	27	6.9	110

Table 4: Overview of West-European monetary (central, low and high) values for the CEN+ indicators

* ES: Ecosystem Services

n.a. = not available; n.r. = not relevant

3 Methodology: selection of data and uncertainties

3.1 Selection of methods and data

Table 5 gives an overview of the available information for monetisation per impact category, taking into account the selected environmental indicators (tables 1 and 2). It distinguishes methods based on damage and prevention costs. The background to the environmental costing methods used is discussed in detail by Debacker et al. (2012).

3.1.1 Indicators for social costs based on damage costs

As damage cost methods calculate in detail how emissions or use of resources causes damages to man and the economy, it is the preferred method. Table 5 illustrates that for most impact categories, there is information on damage costs, although the amount and quality of information differs, which is reflected in the + to +++ scores. We have to look into the details of the available information per impact category to appreciate its strengths and weaknesses. We summarise this info below, a more profound analysis including references is provided in the sections per impact category.

- Damage cost methods calculate how emissions of air pollutants disperse in the atmosphere, affects agriculture and public health which leads to welfare losses in terms of additional costs (e.g. liming, medicine costs), loss of income (production losses, sick leaves) or comfort (e.g. pain). This requires fate and effect models, to account for dispersion, interaction and for impact assessment. The monetary valuation of these impacts uses different methods and data, appropriate to the type of impact. It includes market data (to value loss of agricultural production, medicine costs or loss of income) and data from scientific literature to value pain or loss of life expectancy, based on revealed or stated willingness to pay. Damage costs information is available for most impact categories, and especially if impacts on public health and agriculture are important impact categories (e.g. depletion of the stratospheric ozone layer, tropospheric ozone, human toxicity, particulate matter). A higher number of studies with converging insights is reflected in the scores in table 5 (+++ to ++).
- For impact categories related to depletion of natural resources and with dominant impacts on ecosystems, there is less and more divergent info on damage costs. This is reflected in the + qualification in table 5. Here, it also requires fate and effect modelling to quantify impacts that can be valued. The monetary valuation of impacts on ecosystems for acidification and ecotoxicity is based on stated willingness to pay for biodiversity, (and on prevention costs (see below)). For land occupation, valuation is based on the loss of ecosystem services due to urban land use in Flanders and Europe. The loss of regulatory ecosystem services is valued based on available information per service, either damage cost (air quality, noise) or prevention costs (water quality). The loss of cultural services is based on methods to value health impacts and revealed and stated willingness to pay to live and recreate in a green environment.
- Damages for resource depletion refer to additional costs for future generations, e.g. for mining or importing water over larger distances. These additional costs are based on market prices for e.g. additional energy use.
- For global warming, although there is a lot of information on damage costs, it is qualified with a single +, because this information is less complete and less convergent compared to other impact categories.

Environmental indicator (CEN)	Damage Costs (DC)	Prevention Costs (PC)	Central estimate	Range low-high estimate
1. Global warming	+	++	PC	DC, PC
2. Depletion of the stratospheric ozone layer	++	+	DC	DC, PC
3. Acidification of land and water sources	++	+	DC	DC, PC
4. Eutrophication	+	+	DC, PC	DC, PC
5. Formation of tropospheric ozone photochemical oxidants	+++	-	DC	DC
6. Abiotic depletion of non-fossil resources	+	-	DC	DC, PC
7. Abiotic depletion of fossil resources	+	+	DC	DC, PC
Environmental indicator (CEN+)				
8. Human toxicity, a. cancer b. non-cancer effects	++ +	+ +	DC DC	DC, PC DC, PC
9. Particulate matter	+++	+	DC	DC
10. Ionising radiation, human health	++	+	DC	DC
11. Ecotoxicity: a. terrestrial b. freshwater c. marine		+	DC	
12. Water scarcity	+	-	DC	DC
13. Land use: occupation: a. soil organic matter b. biodiversity b1 : urban b2 : agriculture b3: forestry	+ + + +	+ - + +	DC, PC DC PC PC	DC,PC DC PC PC
14. Land use: transformation a. soil organic matter b. biodiversity b1 : urban b2 : agriculture b3: forestry (excl. rainforest) b4 : rainforest (World estimate)	+ - - - +	+ - - -	DC, PC DC	DC, PC DC

Table 5: Overview of available methods per impact category and methods used for central, low and high estimations

3.1.2 Indicators for social costs based on prevention costs

Prevention cost methods (also referred to as control or abatement costs methods) values an impact based on marginal cost to meet the policy objective for this impact. This requires a clear policy objective (either in terms of emission reductions or environmental quality objectives (such as ppm in ambient air)). In addition, information about all potential prevention measures in different sectors is ranked in terms of their cost-effectiveness, and expressed e.g. in €/kg pollutant. The costs of the least cost-efficient measure to meet the target is an indicator of the value society is willing to pay or impose on citizens or firms to control the environmental problem.

As indicated in table 5, prevention costs are less available, because it requires clear policy objectives and a cost-effectiveness analysis. It is available for global warming and European air quality issues. This is reflected in the ++ in table 5 for global warming, for which a long term

global target is available, which is used in different cost-effectiveness studies and that info is used in several studies and policy documents as an indicator for the social costs of carbon. For impacts related to European air quality (acidification, ozone formation, particulate matter), we qualified the information as + because air quality problem is a multi-source and multi-effect problem which makes it more difficult to assess prevention costs for single effects, because the targets reflect short term compromises rather than long term objectives and because this information is seldom used as indicators for social costs of air pollution.

For impact categories related to ecosystems and biodiversity, there is not enough information for a proper prevention cost method. As a proxy, the costs for a typical measure or the amount of environmental taxes are used. The restoration costs in the context of agri-environmental measures in Europe is taken as an indicator of the value European society puts on safeguarding biodiversity and soil organic matter. For the monetary value of biodiversity, it translates into €/PDF, which is further used to value ecotoxicity and some impacts of acidification or eutrophication. For land use occupation, the info is translated into €/m².

3.1.3 Selection of information for central estimate and low-high estimates

As reflected in table 5, the central estimate for most impact categories is based on damage cost approaches. The estimation of low-high estimates account for uncertainty and information from other sources and methods, including that based on prevention costs. As discussed in these sections, both methods give comparable results.

For global warming, the central estimate is based on prevention costs, because damage costs are highly uncertain and prevention cost information is good.

For soil organic matter, the central estimate is based on both methods. For impacts of land use on biodiversity, and ecotoxicity, we can only provide some monetary indicators on one method with less info and larger uncertainties.

Overall, table 5 reflects that there is more information to value CEN indicators, and that for CEN+ indicators, the availability of information is very diverse.

3.1.4 Comparison with literature

Overall, approaches and data compare rather well with the LCIA assessment of European Energy sector (Alberici, 2014). This study distinguishes between a similar list of 18 impacts and use a similar mix of studies, mainly based on damage costs, except for climate change. The study builds on indicators and data from ReCiPe, ExternE-NEEDS and De Bruyn (2010). Although some indicators are different compared to MMG2014 (e.g. depletion), it builds on the same studies for its monetarisation (e.g. ReCiPe).

3.2 Discounting

The value of future impacts is discounted using a social discount rate. The choice of the discount rate has been subject to many studies and debate, but there is no overall agreement on which discount rate to use (Pearce, 2003; Arrow, 2012). These debates have indicated that discount rates may differ, depending on the objective of the analysis (Goulder, 2012), on the time horizon of the impacts (Arrow, 2012, Ochelen, 2010; Lowe 2008; Commissariat Général du Plan, 2005), ethical issues including equity weighting (Anthoff, 2009) and age weighting (WHO, 2014), uncertainty and risk premiums (Romijn, 2013, Kousky, 2011, van den Bergh, 2007). Consequently, the effect of different discount rates on the overall result has to be reflected in the uncertainty analysis (see below).

For the central estimate, we use a 3 % social discount rate. This discount rate is in line with the recommendations of different governmental institutions and studies that give input to the MMG2014 framework. It is in line with:

- the recommendation of the Flemish government, i.e. a 4 % discount rate to be applied for the first 30 years (within one generation) and a declining discount rate for the rest of the period, that results in a 2 % discount rate for the very long term (Ochelen, 2010) . Such a system results in an average discount rate of 3 %, if we assume an equal yearly impact over a time horizon of 300 years.
- The principle of a declining discount rate is in line with recommendations from the French and UK government, although the latter starts from a lower discount rate (3.5 %) (Commissariat Général du Plan, 2005; Lowe 2008).
- Is close to the recommendation for CBA in the Netherlands, i.e. a base line risk-free discount rate of 2,5 % + a risk premium depending on the type of project (van den Bergh, 2007).
- The discount rate used in ReCiPe, e.g. for the assessment of depletion (Goedkoop, 2013).
- The discount rate used by WHO for the calculations of DALYs.

The discount rate is a very important factor for the assessment of the social costs of global warming, and it is more discussed in detail in that context (see below, paragraph 4.1). Discount rates are also relevant for the quantification and valuation of chronic human health impacts. If monetary valuation is based on the value of loss of DALYs, discounting of future health is already accounted for in the quantification of DALYs using a discount rate of 3 % (WHO, 2014). For valuation of health impacts from emissions of particulate matter, the value of future health impacts is accounted for in the approaches for valuation loss of life expectancy (value of a life of year lost) (Alberini, 2007).

3.3 Uncertainty

For the new environment assessment method (MMG2014) we have updated and improved the MMG2012 approach to deal with the uncertainty.

3.3.1 Background

Uncertainty in LCA, LCIA and impact assessment is determined by a large number of factors (Lloyd, 2008; Huybrechts, 2000). For impact assessment and its valuation, it is important to distinguish between uncertainty due to variation (e.g. different locations of emissions) and uncertainty due to limited understanding of impacts of limits in modelling all the interactions (Webster, 2003, Spadaro 2008, Holland, 2013). Studies that account for a systematic and scientific approach for all sources of uncertainty come up with uncertainty ranges that are very large. In the MMG2012 study we used a systematic approach to assess uncertainty covering all steps in the analysis, following the approach used by Spadaro and Rabl (2008) for assessment of damages from air pollutants. This approach has also been used in the framework of the ALPHA methodology, for assessment of benefits of air pollution for the EC (Holland, 2013). In this context, it has been fully applied within the TUBA framework (Treatment of Uncertainty for Benefit Assessment) in order to test robustness of specific answers to specific questions, i.e. are benefits greater than the costs. We used an estimate of the standard deviation (σ) as the indicator to assess and calculate the uncertainty range. The range reflected a 68 % confidence interval, and was calculated as explained in table 5.

In MMG2012, the standard deviation is assessed for each impact category, and it varies between 3 (for the relative more certain impacts of particulate matter on public health) to 5 (for global warming or biodiversity). This range reflects variation in spatial characteristics, and uncertainty due to limitations in our understanding, models and data. This approach is well justified from a scientific point of view.

Most documents or guidelines that develop indicators and data to assess and monetise environmental impacts however only account for selected sources of variation and/or uncertainty (Lloyd, 2008). For example, the handbook on external costs for transport does discuss the uncertainty related to different impact categories (climate change, particulate matter) but accounts only for variation in two parameters (technologies and location) for the reported values (Ricardo, 2014). The Dutch handbook for shadow prices reports two sets of values, reflecting uncertainty related to the method (damage costs versus prevention costs), but does not account for the uncertainty of the values due to other factors (De Bruyn, 2010). The technical document supporting the US recommendation for the social costs of carbon accounts for variation in two parameters (year of GHG-emission and discount rate), but does not account for the full variation of model uncertainty etc., as done in the more scientific studies (e.g. Tol, 2012). The assessment of damages of industrial activities follows for air pollution the approach of the ALPHA assessment method (Holland, 2013). However, for assessment of uncertainties, it does not include the full range of uncertainty assessments (the TUBA method, as mentioned above) but only accounts for one source of uncertainty, i.e. for valuation of health impacts from air pollutants. Consequently, low and high estimates only differ a factor of 2. Although the technical tools to support the EC air quality policy (particulate matter) allows for a full analysis of uncertainties, the policy documents only report uncertainty due to one parameter (valuation of mortality), with a bandwidth of factor 4 (Holland, 2014).

3.3.2 Treatment of uncertainty in MMG2014

In MMG2014, our approach to uncertainty is closer to how this is dealt with in policy studies and we have adapted our approach in different ways:

- First, we distinguished 'variability' and 'uncertainty'. Variability of the monetary values reflects that emissions and impacts in different locations have a different value, reflecting differences in the physical environment (e.g. average temperature or dominant soil types), and in the socio-economic environment (e.g. number of people exposed to pollution, differences in habits or diets, differences in income and preferences). To account better for this variation, we made distinction between emissions and burdens in Flanders/Belgium, Western Europe and the rest of the world. The data for Western Europe are used as the central data set for the public version of the MMG method.
- Second, we narrowed the confidence interval for the presentation of the results. We used a more narrow band for the low and high estimate, which is more in line with ranges used in documents to support policy analysis. For some impact categories, e.g. eutrophication, we used the central value in different studies to define the low and high estimate. For other impact categories, e.g. global warming, we looked at the bandwidth (BW= high estimate/low estimate) used in other guidelines and policy studies.

We noted that these typical ranges are about half the ranges of the 68 % confidence interval used for the MMG2012. To be more in line with standard practice for policy studies, we adapted our approach to calculate the low and high estimate.

- Third, we have reviewed the assessment of the scientific analysis of uncertainty, because we separated variation and uncertainty and because for some impact categories the indicators/units and LCIA models used have changed.

	MMG2012	MMG2014
	Variation and uncertainty	Uncertainty
Parameter used	σ = standard deviation	BW = Bandwidth, expert judgement $\approx \sigma$ (for some impact categories)
Central value	μ g = expected or central value	μ g = expected or central value
Low estimate	μ g/ σ	μ g / ($\sqrt{\text{BW}}$) $\approx \mu$ g / ($\sigma / 2$)
High estimate	μ g * σ	μ g * ($\sqrt{\text{BW}}$) $\approx \mu$ g * ($\sigma / 2$)
		Spatial Variation
Central value		Western Europe
		Flanders / Belgium
		Rest of the world ***

Table 6: Approach to uncertainty in MMG2012 and MMG2014.

* Depending on σ = standard deviation, determined for each impact.

** Depending on the location of the activity and its impact

*** For emissions in rest of the world, a specific approach to low and high estimate is applied (see text).

The value for Western Europe is the central value, with additional values for Flanders and the rest of the world. The low and high estimate for Western Europe and Flanders are based on the bandwidth around the central estimate, calculated as explained in table 5. Compared to MMG2012, the bandwidth is four times smaller. For emissions in the rest of the world, the bandwidth may be much larger if the impacts vary between countries. Therefore, we apply a broader range. The low estimate may be zero. In general, the damages in Flanders and Western-Europe are higher compared to World average, due to high population density, high environmental pressures on ecosystems and high GDP/capita. Therefore, these damages are representative for the higher estimates at world level. Consequently, the high estimate for world level is based on the maximum of the high estimate for Western Europe and Flanders/Belgium.

Environmental indicator (CEN)	MMG2012M	MMG2014	
	Og (1)	Bandwith (2)	\sqrt{BW} (3)
1. Global warming	5	4	2
2. Depletion of the stratospheric ozone layer	4	4	2
3. Acidification of land and water sources	4	4	2
4. Eutrophication	5	9	3
5. Formation of tropospheric ozone photochemical oxidants	4	4	2
6. Abiotic depletion of non-fossil resources	5	16	4
7. Abiotic depletion of fossil resources	/	16	/
Environmental indicator (CEN)			
8. Human toxicity,			
a. cancer	4	16	4
b. non-cancer effects	4	25	5
9. Particulate matter	4	6,7	2,6
10. Ionising radiation, human health			
	4	9	3
11. Ecotoxicity:			
a. terrestrial			
b. freshwater	4	25	5
c. marine			
12. Water scarcity	/	9	3
13. Land use: occupation:	5		
a. soil organic matter		16	4
b. biodiversity		16	4
14. Land use: transformation	5		
a. soil organic matter		16	4
b. biodiversity		16	4

Table 7: Overview of the uncertainty values for all CEN and CEN+ indicators

- (1) Standard deviation for expected result, assuming log-normal distribution, and used to calculate min. and max estimate within the MMG2012 method
- (2) BW = Bandwidth between low and high estimate = high estimate/low estimate, in case low estimate is zero, bandwidth = standard deviation of MMG2012
- (3) \sqrt{BW} = square root of BW, used to calculate low ($1/\sqrt{BW}$) and high estimates ($* \sqrt{BW}$)

3.4 Comparison with the Dutch environment assessment methods for buildings

In the Netherlands, a similar approach is used to quantify and assess the environmental impacts of buildings and its constituting parts. Although environmental costing is also used as aggregation method, the number of individual LCIA indicators is nevertheless restricted. The development of the Dutch assessment method is performed by SBK (2011). For the definition of the monetary values SBK refers mainly to a study performed by TNO performed in 2006, in order of the Dutch Ministry of Infrastructure and Environment.

The values used for monetary valuation in the Dutch SBK project (2011) are similar to the values for OVAM MMG2014. The SBK data are based on prevention costs. For global warming, ozone depletion, eutrophication, formation of tropospheric ozone and depletion of non-fossil resources, the Dutch SBK values are between the low and high range of OVAM. For global warming, the

SBK value is on the lower boundary, which rather reflects the prevention costs in the short run. The OVAM MMG2014 value is more in line with recent studies, such as the guidelines for monetary valuation of external costs of transport for the EC (Ricardo-AEA, 2014). It should be noted that SBK takes depletion of fossil fuels into account, whereas MMG2014 does not, and one of the arguments is that global warming is a more stringent constraint than depletion.

For acidification, the SBK value is much higher, but it attributes all the prevention costs of reduction of SO₂ emissions to the 'acidification', whereas these costs should only be partly attributed to this problem and partly to the problem of health impacts from secondary particles. The arguments to justify further cuts in SO₂ emissions mainly build on the latter health impacts (e.g. costs benefit studies of EU air quality policies (CAFE)). Attributing prevention costs to both themes, unlike MMG, explains the high value of 4 €/kg SO₂ eq.

3.4.1.1

Environmental Indicator (CEN)	Unit	Method MMG OVAM			SBK
		Central	Low	High	
Global warming	kg CO ₂ eqv.	0.050	0.025	0.10	0.05
Depletion of the stratospheric ozone layer	kg CFC-11 eqv.	49	25	100	30
Acidification of land and water sources	kg SO ₂ eqv.	0.43	0.22	0.88	4
Eutrophication	kg (PO ₄) ³⁻ eqv.	20	6.60	60	9
Formation of tropospheric ozone photochemical oxidants	kg ethene eqv.	0.48	0	6.60	2
Abiotic depletion of non-fossil resources	kg Sb* eqv.	1.56	0	6.23	0.16
Abiotic depletion of fossil resources	MJ, net caloric value	0	0	0.0065	
	kg Sb* eqv.				0.16*

Table 8: Comparison of indicators for monetary valuation in OVAM MMG and Dutch MMG

*Sb: antimony

4 Monetary indicators for CEN indicators

4.1 Global Warming

4.1.1 Monetary value

A. Monetary indicator for global warming, for studies focusing on the comparison of impacts from different building materials or building lines.

Global Warming		Value € / kg CO ₂ eq		
Indicator	Region	Central	Low	High
Full life cycle Fo	Western Europe	0.050	0.025	0.100
	Flanders / Belgium	0.050	0.025	0.100
	Rest of world	0.050	0.025	0.100
More detailed numbers per phase				
Construction	Western Europe	0.045	0.023	0.090
Use Phase	Western Europe	0.110	0.055	0.220
End of life	Western Europe	0.140	0.070	0.280

Table 9: Comparison of indicators for monetary valuation in OVAM MMG and Dutch MMG

Source: VITO 2014, update 2017

B. Monetary indicator for global warming, for assessment of external costs of buildings in cost-benefit analysis (e.g. for comparison with costs of emission reduction measures).

Global Warming		Value € / kg CO ₂ eq		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	0.100	0.050	0.200
	Flanders / Belgium	0.100	0.050	0.200
	Rest of world	0.100	0.050	0.200
More detailed numbers per phase				
Construction	Western Europe	0.045	0.023	0.090
Use Phase	Western Europe	0.110	0.055	0.220
End of life	Western Europe	0.140	0.070	0.280

Table 9: Comparison of indicators for monetary valuation in OVAM MMG and Dutch MMG

Source: VITO 2014

4.1.2 Methods and data used

4.1.2.1 MMG2012

As for MMG2012, the monetary valuation is based on an analysis of prevention costs. The values reflect the costs for the global economy to limit the emission of greenhouse gases to levels that limit global warming to 2°C. The cost assessment is based on a meta-analysis of different models to reach this target (Kuik, 2008) and on other studies that estimate these costs.

The details are described in MMG2012 (Debacker et al., 2012) and De Nocker et al (2010). This estimate is compared to the available information about the expected damages from global warming. The latter is more uncertain, and the damage estimates are incomplete. Nevertheless, the information on damages that can be estimated using the damage cost approach confirms the estimates of the prevention cost approach.

This approach and data are checked against more recent information on damage and prevention costs. In addition, a better procedure to estimate a single value has been developed.

4.1.2.2 Indicator

The indicator (kg CO₂ eq) remains unchanged.

4.1.2.3 Update on literature on social costs of carbon

First, there are a number of new studies or publications on social costs of carbon based on damage costs estimates, including both new original model studies and review of literature (Waldman, 2014; Hope, 2013; Ackerman, 2011; Tol, 2012; Tol, 2011; Tol, 2009). The new estimates confirm the (large) bandwidth of previous studies, with central estimates that range from 7.3 €/ton CO₂eq (range 1 -72 €/ton CO₂eq; based on Fund 3.9 model) (Waldman, 2014; Tol, 2012) to 86 €/ton CO₂eq (Page 09 model, Hope, 2013) (all values converted to euros based on exchange rates of November 2014). A meta-analysis of different studies confirms the large bandwidth, around a central estimate of 30 €/ton CO₂eq (fitted distribution, median) (Tol, 2009). It should be reminded that these studies do not cover all the impacts, and especially those from more extreme global warming scenarios or uncertain events may be important damage categories (Ackerman, 2011; Stern, 2006; Watkiss, 2008).

Second, recent studies based on the prevention costs confirm that the UN objective to limit climate change to a maximum of 2°C can be achieved if worldwide strong measures are taken in all sectors and countries (Edenhofer, 2010; MC Kinsey & Company, 2009; IEA, 2010, Van Vuuren, 2011). It will require big but achievable investments in the coming decades. All studies indicate that the marginal costs of measures will increase over time, in order to remain on a path towards 2°C stabilization towards 2100. There is also a trade-off; fewer measures in the coming decades (and thus lower marginal costs) will imply more costly measures and higher marginal costs in later decades. The range of marginal costs is similar to the range in the study of Kuik (2009). The meta-analysis of Edenhofer (2010) shows the largest range (60 – 550 \$/ton CO₂eq), with lower band for coming decades and higher band for later decades (2060). The report by McKinsey (2012) focusses on measures in short-medium term (towards 2030) with marginal costs of 60 €/ton CO₂eq (for 2030). After 2030, more expensive measures are required with marginal costs up to 100 €/ton CO₂eq.

Thirdly, there are updates guidelines on the use of social costs of carbon. There is an update of the Handbook for External costs of transport for the EC, DG Mobility and Transport (Ricardo-AEA, 2014). This study builds on the same studies as De Nocker (2010) and MMG2012 report (Debacker et al., 2012) and especially on Kuik (2009) for prevention costs of global warming. The study recommends a central value of 90 €/ton CO₂eq for emissions in 2015. The UK has updated its guidelines for social costs, that are based on prevention costs ranging from 79 €/ton CO₂eq for 2015 to 281 £/ton CO₂eq for 2050. The US government recommends a social cost of carbon for use in cost-benefit analysis with a range of 4 € to 53 €/ton CO₂eq (US DOE, 2010). The difference between the US and European recommendations reflects the differences in climate change policies, in line with the prevention cost method.

4.1.2.4 Estimation of the monetary indicator

As explained in MMG2012, the external costs of emissions of greenhouse gases will increase in the coming decades with 6 % per year, reflecting both that marginal costs of meeting GHG-emission targets will become more costly (as the world's economy grows and targets become

tighter) and that marginal damages increase as the world gets warmer. For MMG2012, we recommended to use a starting value of 0.02 euro/kg CO₂eq in 2010 and an annual increase of 6.2 %.

For the MMG2014 method, we developed a better procedure to estimate a single value, based on the following approach:

- The MMG2012 value of 0.02 euro/kg CO₂eq in 2010 with an annual increase of 6.2 %.
- Distribution of emissions of GHG over time, following the attribution of emissions to different phases as in Debacker et al. (2013) (30 % to the production and replacement, 65 % to the use phase and 5 % to end of life).
- Accounting for past inflation, expected future economic growth (based on IPCC scenarios) and discounting (3 %), in line with the assumptions and approach in Ricardo-AEA (2014). This approach is also consistent with how future impacts of current emissions (e.g. chronic health impacts) are dealt with for other impact categories.

This approach results in three different best estimates, depending on the timing of the emissions, and that are represented by the three phases of the life cycle approach:

- Construction : 0.045 € / kg CO₂ eq
- Use phase : 0.110 € / kg CO₂ eq
- End of life: 0.140 € / kg CO₂ eq

For other pollutants and impacts however, there is not enough information to differentiate monetary valuation of emissions or impacts based on their time path. Nevertheless, as for global warming, it is expected that these will also be impacted by e.g. demographic and economic growth. If the objective is to compare impacts, it is not correct to account for these factors for one impact – because info is available – and not for others.

Therefore, we developed different best estimates, depending on the objective of the study, and the level of detail available and assessment of the importance of different life cycle phase.

If the objective is to compare the relative impact of different impact categories, we recommend to use an estimate A, which is best in line with these of other impact categories. This value corresponds to a rounded number, based on the assessment of emissions in the construction phase.

If the objective is to estimate the full environmental costs of buildings, e.g. for use in a cost-benefit analysis, we recommend to use estimate B, as it accounts for more information and is more in line with the guideline values for external costs of GHG in cost-benefit analysis, as explained below.

The rounded best estimate for assessment in cost benefit analysis amounts to 0.1 €/kg CO₂eq or 100 €/ton CO₂eq. This result is similar to the 90 €/ton CO₂eq from the most recent guidelines for external costs of transport for the EC, DG Mobility and Transport (Ricardo-AEA, 2014). The figure in our study is higher because the processes we look at partly occur further in the future, for which higher costs apply. Our figure is also in line with the methodological guidelines for Germany, that recommend a value of 80 €/ton CO₂eq.

In addition, we calculated specific numbers for the construction, use and end of life phase. These impacts are different as we assume these emissions occur in different periods (2013 for construction; 2013-2063 for use phase and after 2063 for end of life). Numbers have been rounded.

4.1.2.5 Spatial variation

The location of the emissions of GHG has no impact on the damages, nor on the prevention costs (as the targets relate to total worldwide emission limits, and assuming least cost approaches). Consequently, there is no need to adapt the monetary value to the location of the emissions.

4.1.2.6 Uncertainty ranges and minimum and maximum estimates

The above mentioned guidelines (Ricardo-AEA, 2014) and UBA (2012) use a bandwidth of a factor 3 and 3.5 between the minimum and maximum value. The latter reflects the span between the minimum and maximum range of the meta analysis of Kuik (2008).

If we use half of the standard deviation, estimated at 5 in MMG2012, the bandwidth would be 6.2. We account for the fact that in recent years there is more consensus between guidelines on what values to use, and for MMG2014, we use a bandwidth of 4.

A bandwidth of 4 between our minimum and maximum estimate means that:

Low estimate = central estimate / $\sqrt{4}$ = central estimate / 2

High estimate = central estimate * $\sqrt{4}$ = central estimate * 2

4.2 Depletion of the stratospheric ozone layer

4.2.1 Monetary value

Depletion of ozone layer		Value € / kg CFC-11 eq.		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	49.1	25	100
	Flanders / Belgium	49.1	25	100
	Rest of world	49.1	25	100

Table 10: Monetary indicator for depletion of the stratospheric ozone layer

Source: VITO 2014

4.2.2 Methods and data used

4.2.2.1 Indicator and new information

The indicator and central values are the same as in MMG2012. They are based on damage costs for health, agriculture and materials (Goedkoop, 2008; De Bruyn, 2010).

4.2.2.2 Spatial variation.

Similar to global warming, there is no need to adapt the monetary value to the location of the emissions.

4.2.2.3 Uncertainty ranges and minimum and maximum estimates.

We apply the general approach, based on standard deviation estimated in MMG2012 ($\sigma = 4$)

This results in a bandwidth of 4 between our minimum and maximum estimate:

Low estimate = central estimate / $\sqrt{4}$ = central estimate / 2

High estimate = central estimate * $\sqrt{4}$ = central estimate * 2

4.3 Acidification of land and water sources

4.3.1 Monetary value

Acidification land water		Value € / kg SO2 eq.		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	0.43	0.22	0.88
	Flanders / Belgium	1.01	0.5	2.02
	Rest of world	0.17	0.00	2.02

Table 11: Monetary indicator for acidification of land and water sources

Source: VITO 2014

4.3.2 Methods and data used

4.3.2.1 Indicator and new information

The indicator and information are the same as in MMG2012, but the new figures account more specifically for spatial variation. Whereas the central figure in MMG2012 is more close to the figure for Flanders; within this latest version the central figure for Western Europe is now lower.

The data are based on the ExternE-method (EC, 2005). It includes damage costs for impacts on building materials and ecosystems, assessed based on the Ecosense model (ExternE-Needs) and impacts on ecosystems (PDF) valued based on restoration costs (Ott, 2006). It does however not include impacts from SO2 on agriculture.

4.3.2.2 Spatial variation

Differences between Western-Europe and Flanders reflect differences in impacts from emissions due to differences in fate modelling (e.g. wind direction and speed), land use, precipitation, building materials used, etc. Data are calculated by the EcoSense-model, as reported for the cases project. Data for Western Europe are based on average EU 27 and data for Flanders are based on Belgium.

World values are based on data for Western Europe, adapted for differences in GDP/capita between Europe and the world average, while accounting for differences in price levels (Purchase Power Parity (PPP) World Bank). Consequently, the value for the rest of the world is 40 % of the value for Western Europe.

4.3.2.3 Uncertainty ranges and minimum and maximum estimates

For Western Europe and Flanders/Belgium, we apply the general approach, based on standard deviation estimated in MMG2012 ($\sigma = 4$). This results in a bandwidth of 4 between our minimum and maximum estimate:

$$\begin{aligned} \text{Low estimate} &= \text{central estimate} / \sqrt{4} = \text{central estimate} / 2 \\ \text{High estimate} &= \text{central estimate} * \sqrt{4} = \text{central estimate} * 2 \end{aligned}$$

For the low and high value for rest of world region we assume that values can be as low as zero and be as high as the maximum value of the Flemish region, where acidification of land and water sources is (still) problematic.

4.4 Eutrophication

4.4.1 Monetary value

Eutrophication		Value € / kg (PO ₄) ³⁻ eqv.		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	20	6.6	60
	Flanders / Belgium	40	9.2	175
	Rest of world	8	0	175

Table 12: Monetary indicator for eutrophication

Source: VITO 2014

4.4.2 Methods and data used

4.4.2.1 Indicator and assessment models

In the MMG2014 method, the indicator for eutrophication (kg (PO₄)₃₋ eqv and LCIA model to assess this indicator (CML 2012) have changed compared to MMG2012 (ReCiPe).

The environmental costing methods and data to monetise the indicator are the same as in MMG2012. It is based on a literature review of studies that use different methods, including 'Willingness to pay' to avoid eutrophication impacts (Gren, 2008), fate and impact modelling to estimate impacts on biodiversity (ExternE-EcoSense) that valued using 'restoration costs' (Ott, 2006) and studies on the 'prevention costs' to meet the objectives for freshwater quality, as required by the European water framework directive (Broekx et al, 2009). The results are expressed in €/kg N, and are calculated to € / kg (PO₄)₃₋ eq. based on characterization factors of CML (1 kg N= 0.42 kg (PO₄)₃₋).

4.4.2.2 Central value, uncertainty ranges and spatial variation

The procedure to estimate central, low and high value is somewhat different compared to other impact categories. As there is a limited number of studies with results that vary widely, we have used these results to determine the low and high value. Assuming a lognormal distribution of the values, we estimate the central value.

The central value for Western-Europe is 20 € / kg (PO₄)₃₋ eq., with a bandwidth of 6.6 to 60. The low value is based on the impact of emissions of N on ecosystems (PDF), which are valued based on restoration costs from Ott (2006). The high estimate is based on the Willingness to pay to avoid eutrophication (Gren, 2008). The bandwidth between the low and high estimate is a factor of 9. This wide range reflects the big uncertainties involved in estimating and valuing eutrophication impacts. This bandwidth does not reflect all uncertainties, e.g. uncertainties related to the use of different indicators (N versus P) and their relative weights as reported in CML-2012, ReCiPe (Goedkoop et al., 2008) and MIRA (2007) is not included.

Assuming a lognormal distribution and the low and high values, the estimate for the central value for Western Europe is 20 (= 60 / 3).

For Flanders, the low value is based on the same source as for Western Europe, but because of spatial variability, the result is 50 % higher. The high cost estimate is based on the prevention costs to meet the objectives of the Water Framework Directive for freshwater in Flanders

(Broekx et al, 2009). These costs are relatively high because the pressures are high in Flanders, and the more expensive measures are required to meet the WFD targets.

For the rest of the world, we adapt the value of Western Europe for differences in GDP per capita (PPP). For the low and high value for rest of world region we assume that values can be as low as zero and be as high as the maximum in Flanders.

4.5 Formation of tropospheric ozone photochemical oxidants

4.5.1 Monetary value

Ozone (low level) formation		Value € / kg ethene eq.		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	0.48	0	6.6
	Flanders / Belgium	3.3	0	6.6
	Rest of world	0.18	0	6.6

Table 13: Monetary indicator for formation of tropospheric ozone photochemical oxidants

Source: VITO 2014

4.5.2 Methods and data used

4.5.2.1 Indicator and assessment model

In the MMG2014 method the indicator for low level ozone formation (kg ethene eqv; (C₂H₄ eqv) and LCIA model to assess this indicator (CML 2012) have changed compared to MMG2012 (i.e. ReCiPe midpoint, expressed in kg NMVOC). The environmental costing methods and data to monetise the indicator are the same as in MMG2012, but we have used a different reasoning to account for variation and uncertainty.

4.5.2.2 Central value, uncertainty ranges and spatial variation

The data are based on the ExternE-method that estimates the impacts of ozone formation, especially on public health (EC, 2005). The fate modelling is based on the Ecosense model (W-Europe) and on the Beleuros model (data for Flanders). The impact assessment and their valuation is for both based on the ExternE method. This method results in different estimates for different ozone precursors (NO_x and NMVOC).

As the process of ozone formation is a complex one, in which ozone precursors (pollutants that contribute to ozone formation (NO_x and NMVOC) can contribute to both ozone formation and decomposition, the marginal impact of different pollutants varies a lot in size, over space and time. Our approach to the low and high values and spatial variation accounts for these differences.

The low values are based on the contribution of emissions of NO_x. This contribution is nil, because in some situations the addressed substances do not lead to photochemical oxidant formation (ozone precursors can have both the effect of formation as defragmentation of photochemical ozone). The high values are based on impacts of ozone precursors with net contribution to ozone formation (NMVOC). This approach is different compared to MMG2012, in which we only used data for NMVOC.

The data for Western Europe are based on data from the ExternE project (EC, 2005), as calculated by Ecosense and reported for the case projects for EU 26 (IER, 2008). The data for Flanders are based on calculations by De Nocker et al (2010). For the rest of the world, values for Western Europe were adapted for differences in GDP/capita (PPP) between Europe and the rest of the world.

To estimate the high value for Western Europe and Flanders, we applied the general approach, based on a bandwidth of 16.

Central estimate = high estimate $\sqrt[4]{}$ = high estimate / 2)

For Europe and for the rest of world, we assume that the high value can be as high as the maximum for Flanders (i.e. 6.6 € / kg ethene eq.).

4.6 Abiotic depletion of non-fossil resources

4.6.1 Monetary value

Depletion of non-fossil resources		Value € / kg Sb* eqv.		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	1.56	0	6.23
	Flanders / Belgium	1.56	0	6.23
	Rest of world	1.56	0	6.23

Table 14: Monetary indicator for abiotic depletion of non-fossil resources

*Sb: antimony

Source: VITO 2014

4.6.2 Methods and data used

4.6.2.1 Indicator and assessment model

The methods and data to monetise abiotic depletion of non-fossil resources are based on ReCiPe, as in MMG2012. However, as in ReCiPe the indicator is based on \$ / kg Fe eq., and as there is no conversion factor given for Sb, we have to add an additional step to express the costs in terms of €/kg Sb eq. In addition, we have used a different reasoning to account for variation and uncertainty.

4.6.2.2 Central value, uncertainty ranges and spatial variation

Low estimate

The low value is nil, and reflects the point of view that resource depletion costs are internalized in prices (De Bruyn, 2010). This argument is further discussed in MMG2012.

High estimate

The high estimate is based on the additional costs for future generations for extraction of raw materials. The assessment and valuation methods are discussed in MMG2012. It is based on

ReCiPe (Goedkoop, 2008) and the additional costs of energy required for extraction that can be interpreted as a 'resource depletion cost'. It amounts to 0.0518 €/kg Fe, which corresponds to 0.1 % of the market price of Fe (market price of 1 kg Fe equivalent = 72 \$/kg).

The resource depletion costs differ between resources, depending on e.g. the current grade of metals or minerals in the ore. In ReCiPe, the resource depletion costs are calculated for 20 substances. Expressed as a % of the market price of the substances, the resource depletion cost varies from 0.1 % to + 600 %, with a median value of 3.4 % and an average of 88 % (see table below).

ReCiPe does not give a value for Sb. If we were to apply the characterisation factor from ILCD (1 kg Fe= 1.66 E-6 Sb), the resource depletion cost of Sb would amount to 42.2 k€/kg Sb, which is more than 4000 times the market price. If we were to use this characterisation factor as a basis for assessment of resource depletion costs, we would on average overestimate the resource depletion costs with many orders of magnitude (see table 12, method 1, columns 4 to 6). From the 13 substances for which we can make a direct comparison with ReCiPe, only for Fe, Cu and Mn this resource depletion cost estimate is in the same order of magnitude compared to those of ReCiPe. For all other materials, this approach would - compared to ReCiPe - overestimate the resource depletion costs with on average 4 orders of magnitude. If we were to use this approach, the average resource depletion costs would amount to 384 times the market prices. This cannot be justified based on the calculations from ReCiPe.

Therefore, we estimate the resource depletion costs for Sb based on the average value for the ratio between resource depletion costs - as estimated in ReCiPe - and the market price. On average, this ratio is 82.9 % (column (3) of table 15 and 16). At a market price of 9.4 \$/kg Sb, this results in a resource depletion cost of 7.8 \$/kg Sb or 6.23 €/kg Sb. This approach results in resource depletion costs for the different substances that are on average within the same order of magnitude compared to those of ReCiPe. For the majority of the substances, this estimate is lower compared to the ReCiPe estimate.

To estimate the central value, we apply the general approach, based on the standard deviation estimated in MMG2012 ($\sigma = 5$). We use a bandwidth of 16 between the low and high estimate. Thus we calculate:

- Central estimate = high estimate / $\sqrt{16}$ = high estimate / 4
- Central estimate = 6.23 €/kg Sb / 4 = 1.56 €/kg Sb

As the market for raw materials are global markets and depletion is a world-wide problem, we do not apply different values for different countries or regions.

	RDC ReCiPe \$/kg	market price \$/kg	ratio =(1)/(2) %	RDC Method 1 \$/kg	ratio =(4)/(1) %	ratio =(4)/(2) %
	(1)	(2)	(3)	(4)	(5)	(6)
Ag	20,48					
Al	0,0064	1,3	0,49%	1,07	16670%	82%
Ay	5,006					
Co	0,072	46	0,16%	1079,5	1499331%	2347%
Cr	1,78	9,6	19%	0,83	46%	8,6%
Cu	3,06	7	44%	105,4	3445%	1506%
Fe	0,07	72	0,10%	0,07	100%	0,10%
Ir	6,62	42000	0,02%	560,8	8472%	1,3%
Mn	5,48	1,5	365%	0,99	18%	66%
Mo	14,85	30	50%	2998,19	20190%	9994%
Ni	0,9	40	2,25%	176,3	19585%	441%
Os	464	77000	0,60%			
Pb	0,13	2,5	5,20%	632,5	486562%	25301%
Pd	273	58330	0,47%	394699	144578%	677%
Pt	11652	5400	216%	383313	3290%	7098%
Rh	1455	80000	1,82%			
Ru	144	14000	1,03%			
Sn	90,99	13	700%	4849,4	5330%	37303%
Zn	0,16	3,5	4,57%	153,9	96197%	4398%
Sb	na	9,4		42.168,7	na	448603%
Average			82,9%		164.558%	38.416%

Table 15: Resource depletion costs (RDC) for different substances, based on characterization factor ILCD (method 1)

na = not available

(1) RDC = Resource depletion costs, as estimated in ReCiPe (Goedkoop, 2013) p 117, in \$/kg

(2) Market prices, in \$/kg (Wikipedia 2014, compiled from different sources)

(4) RDC = Resource depletion costs, using method 1 = based on the RDC for Fe, the characterization factors from ILCD for Sb (1 kg Fe= 1.66 E-6 Sb) and for other metals.

	RDC ReCiPe \$/kg	market price \$/kg	ratio =(1)/(2) %	RDC Method 2 \$/kg	ratio =(4)/(1) %	ratio =(4)/(2) %
	(1)	(2)	(3)	(4)	(5)	(6)
Al	0,0064	1,3	0,49%	0,0002	3,1%	0,02%
Co	0,072	46	0,16%	0,200	277%	0,43%
Cr	1,78	9,6	19%	0,0002	0,0%	0,00%
Cu	3,06	7	44%	0,019	0,6%	0,28%
Fe	0,07	72	0,10%	0,00001	0,0%	0,00%
Ir	6,62	42000	0,02%	0,10	1,6%	0,00%
Mn	5,48	1,5	365%	0,0002	0,0%	0,01%
Mo	14,85	30	50%	0,55	3,7%	1,85%
Ni	0,9	40	2,25%	0,033	3,6%	0,08%
Os	464	77000	0,60%			
Pb	0,13	2,5	5,20%	0,117	89,9%	4,68%
Pd	273	58330	0,47%	72,95	26,7%	0,13%
Pt	11652	5400	216%	70,84	0,6%	1,31%
Rh	1455	80000	1,82%			
Ru	144	14000	1,03%			
Sn	90,99	13	700%	0,896	1,0%	6,89%
Zn	0,16	3,5	4,57%	0,028	17,8%	0,81%
Sb	na	9,4		7,8		82,9%
Average			82,9%	10,24	30,4%	6,6%

Table 16: Resource depletion costs (RDC) for different substances, based on ratio to market prices (method 2)

- (1) RDC = Resource depletion costs, as estimated in ReCiPe, Goedkoop, 2013, p 117, in \$/kg
(2) Market prices, in \$/kg (Wikipedia 2014, compiled from different sources)
(4) RDC = Resource depletion costs, using method 2 = based on the average ratio RDC/market price and applied to Sb market prices for Fe, the characterization factors from ILDC for Sb (1 kg Fe= 1,66 E-6 Sb) and for other metals.

4.7 Abiotic depletion of fossil resources

4.7.1 Monetary value

Depletion of fossil resources		Value € / MJ, net caloric value		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	0	0	0.0065
	Flanders / Belgium	0	0	0.0065
	Rest of world	0	0	0.0065

Table 17: Monetary indicator for abiotic depletion of fossil resources

Source: VITO 2014

4.7.2 Methods and data used

4.7.2.1 Indicator and assessment model

As in the MMG2012 method our best estimate for the central value is a zero value (Debacker et al., 2012), we have added now a value for the high estimate.

4.7.2.2 Central value, uncertainty ranges and spatial variation

The central and the low value is zero. This reflects the point of view that resource depletion costs are internalized in market prices (De Bruyn, 2010, Aguilera 2009, Aguilera 2013). In addition, MMG2014 uses a valuation of greenhouse gas emissions based on prevention costs, which assume that we limit emissions of greenhouse gasses to limit global warming to a maximum of 2°C, in line with the UN objective. This emission path limits the use of fossil fuels, irrespective of its availability (De Nocker, 2010).

The high value is based on the Eco-indicator 99 method. (Allacker et al, 2012) and costs of military action to secure energy supply (NDCF, 2007; Stern, 2010 as cited in VTPI, 2013).

As energy markets are global markets and depletion is a world-wide problem, we do not apply different values for different countries or regions.

5 Monetary indicators for CEN+ indicators

5.1 Human toxicity, cancer effects

5.1.1 Monetary value

Human toxicity cancer		€/ CTUh*		
	Region	Central	Low	High
	Western Europe	665,109	166,277	2,660,434
	Flanders/Belgium	680,451	170,113	2,721,802
	rest of world	266,043	66,511	2,721,802

Table 18: Monetary indicator for human toxicity cancer

*CTUh = (comparative toxic units human health)

Source: VITO 2014

5.1.2 Methods and data used

In the MMG2012 method, impacts on human health are assessed based on ReCiPe (Goedkoop et al., 2008), and expressed in terms of DALY (disability adjusted life years).

In the MMG2014 method, the impacts are assessed based on the USEtox method (Rosenbaum, 2008) and expressed in CTUh (comparative toxic units human health).

Quantification: The CTUh case = 1 cancer = 11.5 DALY (Rosenbaum, 2008; JRC 2011, p 31.)

Valuation :

Costs of cancer = cost of medical care (a) + loss of production value (b) + loss of life expectancy (c)

(a) + (b) are estimated based on a study of total costs of cancer in EU = 51,429.6 € (Luengo-Fernandez, 2013)

(c) Assuming 1 DALY related to cancer corresponds to 1 YOLL (years of life lost), then the value of a life year lost (VOLY) = 53,363.5 €/DALY (W-Europe)

CTUh cancer = 51,429.6 € + (11.5 x 53,363.5) = 665,109 € / CTUh.

5.1.3 Further explanation

The assessment is based on the scientific consensus model USEtox, developed in the framework of the UNEP–SETAC Life Cycle Initiative (Rosenbaum, 2008).

The characterisation factor for human toxicity (human toxicity potential) is expressed in comparative toxic units (CTUh). CTUh describes the estimated increase in morbidity in the total human population per unit mass of a chemical emitted (cases per kilogram), assuming equal weighting between cancer and non-cancer due to a lack of more precise insights into this issue (Rosenbaum, 2008). The cases can be converted to DALYs based on information of the ILCD Handbook (2011), making a distinction between cancer and non-cancer cases.

The total costs of a cancer case include:

- (a) health care costs;
- (b) loss of productivity for patient and family;
- (c) loss of value of life expectancy for the patient.

The costs related to health care (a) + loss of productivity (b)

- These costs are quantified based on a recent study of total costs of cancer for the EU published in the Lancet Oncology (Luengo-Fernandez, 2013). It estimates total costs in the EU to be 126 billion (2009) of which 40 % for health care costs. These total costs per case of cancer (diagnosed) is 51,429 euro.
- This figure is substantially lower than the 450,000 euro/cancer case used in ExternE, but the sources for these latter data are not well documented (Eftec, 2004). Health care costs for the EU given by Luengo-Fernandez (2013) are in the same order of magnitude as health care costs for the US (Yabroff, 2007 cited in Mariotto, 2011).
- We assume the same value for Western Europe and Flanders. Despite the similarities between Europe and the US, we adapted the value for the rest of the world based on the differences in GDP/capita (PPP) between Europe and world.

The costs related to loss of value of life expectancy for the patient (c)

- In addition, it is estimated that on average a cancer case results in 11.5 DALYs loss of years of health life (JRC 2011), based on Huijbregts (2005). We assume that the value of 1 DALY for cancer equals 1 YOLL (year of life lost). It has been discussed in literature to which extent one should apply a 'cancer premium'. Some studies have found such premium which reflects a 'dread' factor and accounts for more pain and suffering associated with cancer (Eftec, 2004; Cropper et al, 2011). We do not apply such a cancer premium, because the evidence in literature is mixed. Whereas e.g. Hammit (2004) and Van Houtven (2008) found an important cancer premium (at least a factor 1.3), other recent studies Hammit and Haninger (2010) Adamowicz et al. (2009) did not.

Valuation of a DALY:

We assume that the value of 1 DALY (disability adjusted life year) for cancer equals 1 YOLL (year of life lost), which is well justified for valuation of health impacts related to cancer. It has also been demonstrated that for the valuation of the major morbidity impacts the results using DALYs or using other data from economic literature results in identical impacts (Desaigues, 2007).

For the valuation of a YOLL, we use the figures for EU-25, based on survey in different EU countries in the context of the EU Needs project (40000 € in €2000 prices) (Desaigues et al, 2011) and corrected for inflation to 2012 price levels based on Eurostat HICP data).

- Western Europe = 1 DALY = 1 YOLL = 53,363 € (based on values for Europe in Desaigues et al, 2011)
- Valuation data for Flanders are based on EU wide data. As values depend on income per capita, the NEEDS study recommends to adapt the figures for local GDP/capita. GDP/capita in Flanders is 2.5 % above EU-average. 1 DALY = 54,697 €
- Rest of World: 1 DALY = 1 YOLL = 21,345 € (=YOLL (based on EU2013 value and differences in GDP/capita (PPP) between Europe and world).

For comparison: the economic valuation of a DALY accounting for all information on mortality and morbidity gives much broader ranges (from 5,000 € to 400,000 €). Central estimates vary from 10.000 to 100,000 €.

5.1.4 Low and high values

Rosenbaum reports uncertainty boundaries that reflect a standard deviation of 20 (Rosenbaum, 2008 and JRC, 2011). We have no yardsticks to compare this uncertainty assessment with that of other impact categories. The large uncertainty boundaries for human toxicity reflects that this impact category involves a wide range of exposure routes for a wide range of chemicals, and that for a large number of these chemicals the information of potential effects on human health is scarce, and has to be estimated based on e.g. animal experiments. In this context, it is common to work with uncertainty boundaries covering several orders of magnitude (Rosenbaum, 2008). A particular issue is that for a number of substances, information is missing, and impacts are estimated as zero impacts (JRC, 2011). Since the publishing of the assessment of USEtox method within the framework of the ILCD Handbook (2011) more chemical substances have been modelled.

Because impacts are both relatively uncertain and highly variable, we use a relative high bandwidth of 16.

Low value = central value / $\sqrt{16}$ = central value / 4
 High value = central value * $\sqrt{16}$ = central value * 4

The estimate for Flanders and for the rest of the world are based on the figure for Western Europe, adapted for differences in GDP per capita. For rest of the world, we assume that the maximum value = max value for Flanders.

5.2 Human toxicity, non-cancer effects

5.2.1 Monetary value

Human toxicity non-cancer		€/ CTUh*		
	Region	Central	Low	High
	Western Europe	144.081	28.816	720.407
	Flanders/Belgium	147.683	29.537	738.417
	rest of world	57.633	11.527	738.417

Table 19: Monetary indicator for human toxicity non-cancer

* CTUh = comparative toxic units human health

Source: VITO 2014

5.2.2 Methods and data used

The environmental costing method is similar to that for human toxicity cancer. The main difference is the interpretation of a CTUh for non-cancer.

Quantification: 1 CTUh case = 1 non-cancer = 2.7 DALY (Rosenbaum, 2008; JRC, 2011, p 31)
 The valuation of DALY is discussed above.

The uncertainties are larger for non-cancer compared to cancer (Rosenbaum, 2008 and JRC, 2011). To reflect this difference, we assume a bandwidth of 25 between the low and high value:

Low value = central value / $\sqrt{25}$ = central value / 5
 High value = central value * $\sqrt{25}$ = central value * 5

5.3 Particulate matter

5.3.1 Monetary value

Particulate matter		Value € / kg PM 2.5		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	34.0	12.7	85.0
	Flanders / Belgium	57.8	21.7	144.5
	Rest of world	7.7	2.9	19.2

Table 20: Monetary indicator for Particulate matter

Source: VITO 2014

5.3.2 Methods and data used

In the MMG2012 assessment method, assessment and valuation were based on ReCiPe (Goedkoop et al., 2008). The data were based on the application of the ExternE accounting framework (Preiss and Klotz, 2008) to Flanders, in the context of a project for VMM and using specific air quality models and data for Flanders (De Nocker, 2010). In the MMG2014 assessment method, environmental costing models are also based on the ExternE accounting framework (Preiss and Klotz, 2008), as applied to EU 28 countries, using specific air quality models and data for EU 28.

Because the indicator and LCIA model have changed, we explain more in detail how the values were calculated. This level of detail is also justified because impacts of PM on public health are likely to be an important part of total monetized impacts for most applications.

Step 1: fate, exposure and effect analysis

- We use the results from the European LC-Impact project (2010-2013) (Preiss, 2013). In the LC-Impact project, estimates for primary PM and for PM precursors have been estimated for emissions from all regions (countries) in the world, building on earlier work and using the models that have been identified by ILCD as the best to use (RiskPoll, Humbert) (Preiss, 2013, ILCD, 2008). Emissions of precursors (SO₂, NO_x and NH₃) will lead to an increase of PM aerosols in ambient air, and to related health impacts.
- The main progress of the LCA-Impact model relates to the impact factor, estimating the impact of emissions on intake by humans, while accounting for differences in population density exposed to the emission, distance between emission and population exposed (typically stack height) and meteorological conditions, especially wind speed and atmospheric mixing height. These differences are dealt with using ‘archetypes’ (remote, rural, urban) for emissions, as developed by Humbert (2011) (and reflecting the work of the UNEP/SETAC) Life Cycle Initiative). The impacts are modelled combining several dispersion models.
- The effect factor builds on the dose-response functions as developed in the EU-Heimsta project (Torfs et al, 2007) used for many assessments by the EU and WHO, and their aggregation into DALYs (as defined by WHO, 2012). The effect estimates account for differences in toxicity between primary PM and PM precursors and differences between countries related to age distribution.
- As final deliverables were not yet available, we used the results of the project as presented at SETAC 2013. European values are based on the average of values for Western and Eastern Europe (although differences are small). Impacts for emissions in Flanders are higher compared to EU-average (on average 1.8 times). This is based on the relationship

between estimates for Belgium and EU average, as assessed by LC-Impact (PM (average for low and high stacks) or Cases (SO₂, NO_x, NH₃ (unspecified stack heights)).

—
Step 2: valuation

- As the major impact from particulate matter relates to impacts on mortality (loss of life expectancy), we use the information on valuation of mortality impacts from PM to value a DALY in monetary terms. This valuation is based on the value of a YOLL (Year of life lost). We assume that 1 DALY equals 1 YOLL, which is well justified for valuation of health impacts from PM. It has also been demonstrated that for the valuation of the major morbidity impacts the results using DALYs or using other data from economic literature results in identical impacts (Desaigues, 2007).
- For the valuation of a YOLL, we use the figures for EU-25, based on survey in different EU countries in the context of the EU Needs project (40000 € in €2000 prices) (Desaigues et al, 2011) and corrected for inflation to 2012 price levels based on Eurostat HICP data.
 - Western Europe: 1 DALY = 1 YOLL = 53.363 €
 - Valuation data for Flanders are based on EU wide data. As values depend on income per capita, the NEEDS study recommends to adapt the figures for local GDP/capita. GDP/capita in Flanders is 2.5 % above EU-average. 1 DALY = 54.697 €
 - Rest of the world: 1 DALY = 21.345 €/=YOLL (based on EU2013 value and differences in GDP/capita (PPP) between Europe and world).

For comparison: the economic valuation of a DALY accounting for all information on mortality and morbidity gives much broader ranges (from 5.000€ to 400.000€. Central estimates vary from 10.000 to 100.000 €).

Results: step 1 x step 2

See table 21 to table 24

5.3.3 Low and high values

The uncertainty boundaries for PM 2.5 especially reflect a wide variation in impacts, depending on variations in fate and exposure assessment.

There is a wide variety in impacts and effects, depending on the variation in situations (population density, stack height, etc.). The variation is very high (two orders of magnitude) for emissions of different sectors and countries, but there is less information about the variation if applied in LCIA analysis, where emissions are always a combination of emissions from different countries, regions and/or sectors.

Although uncertainties remain related to impact assessment and their valuation, there is more consensus and certainty about the impacts per kg inhaled, as these elements have been subject to many studies and policy decisions over the last 20 years. This is e.g. reflected in the fact that most assessment models use similar dose-response functions, at least for the most important impacts.

To calculate low and high values, we have applied an uncertainty range of a factor of 2 for impact assessment, which reflects the differences between models (Preiss, 2013). In addition, we have used a range of +/- 25 % to account for variation and uncertainty in valuation. If we apply these ranges to the estimates of DALY/ton and value/DALY, it results in a bandwidth of 6.7 between the high and low estimate.

This is a wider range compared to most other impact categories. It does not mean that these impacts are more uncertain, but that there is a larger variation in impact between sites

(emissions in rural versus urban areas) and between sectors (high stack emissions versus low stack emissions (transport)).

Discussion:

For PM_{2.5}, the results relate very well to the estimates based on the more detailed EcoSense runs, as used in and reported for the EU CASES project (Feem, 2008). For SO₂, NO_x and NH₃, LC-Impact results are lower compared to LC-Impact, but fall within the uncertainty boundaries as calculated in table 21.

It has to be noted that the uncertainty boundaries are similar to those for global warming, although the rationale behind is rather the opposite. For global warming, there is a consensus and little uncertainty about fate and exposure for different GHG emissions from different sectors and countries (contribution to global warming), but large uncertainties related to the impacts and valuation of global warming. For PM, there is a lot of consensus about effects and their valuation, but large variation in fate and exposure for emissions from different sectors and countries.

Particulate matter				
€/PM 2.5 kg	Region	Central	Low	High
	Western Europe	34.0	12.7	85.0
	Flanders	57.8	21.7	144.5
	World	7.7	2.9	19.2
€/NO _x kg	Region	Central	Low	High
	Western Europe	5.1	1.9	12.8
	Flanders	7.2	2.7	17.9
	World	1.3	0.5	3.3
€/SO ₂ kg	Region	Central	Low	High
	Western Europe	5.5	2.1	13.7
	Flanders	9.0	3.4	22.4
	World	2.9	1.1	7.3
€/NH ₃ kg	Region	Central	Low	High
	Western Europe	17.2	6.4	42.9
	Flanders	47.0	17.6	117.4
	World	2.6	1.0	6.6

Table 21: Damage costs per kg emitted, for primary particulate matter and particulate matter precursors

Particulate matter : background information, impact assessment (DALY/ ton emitted)				
DALY/PM 2.5 ton	Region	Central	Low	High
	Western Europe	0.64	0.32	1.27
	Flanders	1.06	0.53	2.11
	World	0.40	0.20	0.79
DALY/ NO _x ton	Region	Central	Low	High
	Western Europe	0.10	0.05	0.19
	Flanders	0.13	0.07	0.26
	World	0.07	0.03	0.14
DALY/SO ₂ ton	Region	Central	Low	High
	Western Europe	0.10	0.05	0.21
	Flanders	0.16	0.08	0.33
	World	0.15	0.08	0.30
DALY/NH ₃ ton	Region	Central	Low	High
	Western Europe	0.32	0.16	0.64
	Flanders	0.86	0.43	1.72
	World	0.14	0.07	0.27

Table 22: Impacts per ton emitted, for primary particulate matter and particulate matter precursors

Method: damage based: Data sources: LC-Impact (Preis, 2013)

Valuation of DALY	background information, valuation (€/ DALY)		
Region	Central	Low	High
Western Europe	53,363	40,023	66,704
Flanders	54,698	41,023	68,372
World	19,816	6,605	68,372

Table 23: Valuation of health impacts, DALY (disability adjusted life years)

Low (high) values = central value - / + 0.25 * central value

External costs		In € / kg	
Pollutant	Sector	EU 27	Belgium
PM 2.5	Unspecified sector and stack height	31.2	60.9
	Low stacks (households)	32.8	61.4
	High stacks (power, industry)	15.5	26.3

Table 24: Impacts from emissions calculated by EcoSense: for year 2015, in EU 27 and Belgium

Source: Cases, FEEM, 2008

5.4 Ionising radiation: human health effects

5.4.1 Monetary value

Ionising radiation, human health		Value € / kg U235 eqv.		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	9.7E-04	3.2E-04	2.9E-03
	Flanders / Belgium	9.9E-04	3.3E-04	3.0E-03
	Rest of world	3.6E-04	1.2E-04	3.0E-03

Table 25: Monetary indicator for ionising radiation, human health

Source: VITO 2014

5.4.2 Methods and data used

In the MMG2012 assessment method the fate and impact assessment was based on the endpoint methods within ReCiPe. For valuation, the monetary indicator for DALY was used.

In the MMG2014 assessment method, quantification is based on midpoint assessment (Frishknecht et al, 2000) as implemented in ReCiPe midpoint assessment.

Quantification

The exposure and fate factor for 1 kg of uranium U235 eqv. is 1.40E-08 Man Sievert (Man.Sv) per kBq (Goedkoop et al. 2008, Table 9.1).

Second, the impact factor per man.Sv is 1.17 DALYs (Goedkoop et al. 2008, Table 9.3).

The total number of DALYs per kg U235 eq. is 1.64E-08.

Valuation

The valuation of a DALY in this context needs to account for the type of health impacts. The health impacts from exposure to ionising radiation relate to cancers (fatal and non-fatal) and severe hereditary effects.

For hereditary effects, the loss of life expectancy is estimated at 59 years (Goedkoop et al. 2008), and – in line with valuation of life expectancy for other health impacts – loss of 1 year is estimated at 53 k€. The valuation of cancer accounts - As explained above (paragraph 5.8) - for loss of life expectancy (part A) and in addition the costs of medicine and loss of labour income (part B). These costs amount to 51 k€ (as explained in paragraph 5.8), divided by the number of DALYs, i.e. 11.5 for fatal cancers and 2.7 for non-fatal cancers.

Accounting for the relative share of these endpoints in the total numbers of DALYs per man.Sv, the average value per DALY is 69,305 €.

Consequently, the value per kg U235 eqv. is 9.7E-04 (central estimate for Western Europe).

	Quantification DALYs /mSv	Valuation €/DALY			Valuation €/mSv	
		Part a life exp. (2)	Part b medicine (3)	Total (4)	Total (5)	%
	(1)					
Fatal cancers	0.45	53,363	4,472	57,836	26,131	38%
Non-fatal cancers	0.25	53,363	19,048	72,411	18,435	27%
Severe hereditary effects	0.46	53,363	0	53,363	24,739	36%
Total per m.Sv	1.17				69,305	100%

Table 26: Valuation of DALY for ionising radiation, for Western Europe

(1) Goedkoop et al., 2008, p. 85

(2) Costs related to loss of life expectancy (Assuming 1 DALY = 1 VOLY, valuation, see table 23, for W-Europe)

(3) Costs related to health care and loss of productivity (51 €/11.7 DALYs for fatal cancers and 2.7 DALYs for non-fatal cancers.

(4) (4)= (2) + (3)

(5) (5)= (1) x (4) t.

5.4.2.1 Uncertainty and low and high estimates

We use a bandwidth of 9, which is larger than for e.g. particulate matter and that reflects that the valuation of severe hereditary effects based on DALYs is very uncertain. It is narrower than for human toxicity, because for ionising radiation the impacts are associated with a limited number of substances.

The estimates for Flanders and for the rest of the world are based on the figure for Western Europe, adapted for differences in GDP per capita. For rest of the world, we assume that the maximum value equals the max value for Flanders.

5.5 Ecotoxicity, freshwater

5.5.1 Monetary value

Ecotoxicity, freshwater		Value €/CTUe*		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	3.70E-05	7.39E-06	1.85E-04
	Flanders / Belgium	3.70E-05	7.39E-06	1.85E-04
	Rest of world	1.34E-05	2.68E-06	1.85E-04

Table 27: Monetary indicator for Ecotoxicity, freshwater

* CTUe = comparative toxic unit for ecosystems

Source: VITO 2014

5.5.2 Methods and data used

In the MMG2012 assessment method the fate and impact assessment was based on the endpoint method ReCiPe, expressed in PDF/m² (potentially disappeared fraction) and PDF was valued based on restoration costs (Ott, 2006). The indicator used for quantification was expressed in kg 1.4 dichlorobenzene (DB) equivalent.

In the MMG2014, assessment method the characterization factor for aquatic ecotoxicity (ecotoxicity potential) is expressed in comparative toxic units for ecosystems (CTUe) and provides an estimate of the potentially fraction of affected species (PAF) integrated over time and volume per unit mass of a chemical emitted (PAF m³ day kg⁻¹) (Rosenbaum, 2008, p. 7).

Quantification: 1 CTUe, freshwater = 1 PAF/m³ day/kg.

Valuation

We use the value €/PDF as discussed in the MMG2012 assessment method, i.e. 0.46 €/PDF. This value corresponds to the value attached to protection of terrestrial biodiversity, and is based on the costs citizens say are willing to pay to protect biodiversity. The underlying assumption is that all terrestrial species are valued equally.

To value ecotoxicity in freshwater, we have to expand the assumption, and assume that freshwater species are valued identical to terrestrial species. In addition, we assume that the values discussed above relate to all terrestrial species, and that aquatic species are valued on top of these, and that the value as discussed above can be used for aquatic species. This corresponds to 3.7 E-05 €/CTUe. The details of the calculation are given in Table 29.

Indicators	correction factor	Value
Economic value of terrestrial biodiversity in PDF m ² .year/kg		0,46
from PDF m ² terrestrial to PDF m ³ freshwater (1)	0,053	0,025
from PDF to PAF (2)	0,55	0,013
from PAF year to PAF day (3)	0,00273973	0,0000370
Value ecotoxicity , €/CTUe (in PAF.m ³ .day/kg)		3,70E-05

Table 28: Correction factors used for valuation of Ecotoxicity, freshwater

- (1) Assuming equal value of terrestrial and freshwater species, number of species based on Goedkoop, 2008, update 2011
- (2) Based on Goedkoop, 2008 (based on Van Zelm 2007 and 2009)
- (3) Accounting for 365 days/year

5.5.2.1 Uncertainty and low and high estimates

Given the wide range of species and substances involved, the uncertainty and variation for assessment and valuation of ecotoxicity is at least as great as that for human toxicity cancer. In addition, valuation of impacts on biodiversity is much more uncertain compared to that of human health. Consequently, the uncertainty range is at least as high as that for human-health non-cancer. This explains the bandwidth of 25 to express this relative large uncertainty and variation.

Low value = central value / $\sqrt{25}$ = central value / 5
 High value = central value * $\sqrt{25}$ = central value * 5

5.6 Water scarcity

5.6.1 Monetary value

Water scarcity		Value € / m ³ water eqv.		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	0.067	0.022	0.200
	Flanders / Belgium	0.079	0.026	0.238
	Rest of world	0.012	0.004	0.238

Table 29: Monetary indicator for Water use

Source: VITO 2014

5.6.2 Methods and data used

Introduction

In the MMG2012 assessment method, this impact category was not monetized. Therefore, we give more details about the calculation of the monetary values in the current assessment method.

As water is a renewable source, the environmental and socio-economic impacts of the use of water within the limits of the resource availability, are very limited. The costs of water use above these limits may be very high, depending on the value of the foregone water use. The impacts may be environmental (low water impacts in rivers and drought impacts on habitats) but also socio-economic, e.g. limiting the use of water in other sectors that may add more value to a m³ of water compared to the current uses.

From a socio-economic perspective, the use of water is very different compared to e.g. the use of mineral resources or fossil energy sources. Whereas the economist may argue that scarcity of mineral and fossil resources will be reflected in prices and markets, this is not the case for the use of water because there are no real markets for water. The right to use water and the prices paid for it depend on (often complex) historical rights and specific institutional settings and agreements. In general, the price of water that users pay reflects – at maximum – the financing investments and operational costs of water treatment and water distribution systems, but not for the use of the water resource itself. As a consequence, the institutional arrangements (e.g. permits) and the price mechanisms do not reflect water scarcity. On the other hand, there is a lot of evidence that water users do not always pay the full financial costs and water use is subsidized (FAO, 2004; Anderson, 2008). This will lead to inefficient use of water, and limits the use of water savings or alternative water sources.

Water scarcity is a local, and sometimes temporary problem, which is difficult to deal with in generic terms. Furthermore, the understanding of the issue, indicators to measure water scarcity or legal frameworks to deal with it are less far developed compared to water quality (EC, 2007). As a consequence, LCIA methods have not much indicators, standards or criteria to build upon.

Step 1: fate, exposure and effect analysis

Compared to e.g. particulate matter, there is no history of complex models used to account in detail for variation in water scarcity. We use the Swiss Ecoscarcity method, as this method offers on the one hand some regional differentiation, and is well accepted (Kounina, 2013). It has

however to be noted that more recently other methods have been developed, that go a step further in terms of a more detailed assessment of water use and accounting for regional differences in water scarcity (Kounina, 2013).

The Swiss Ecoscarcity method accounts for the water scarcity in the region where water is used. To this purpose, it calculates a weighting factor that is equal to: (Frischknecht, 2006)

$$\text{weighting factor} = \left(\frac{\text{Water consumed in a region A (m}^3\text{)}}{(\text{Renewable water resource in region A (m}^3\text{)}) \times (\text{rate of sustainable use (\%)})} \right)^2$$

It further assumes the rate of sustainable use to be 20 %, based on OECD that states that regions with 20 % use of renewable water sources have a medium water stress (Frischknecht, 2006). This % is in line with criteria used by the European Environmental Agency (EEA, 2009)

The weighting factor varies from 0.0625 for countries or regions with a "low water pressure" (e.g. Switzerland) to 2.25 for countries with a "medium water pressure" (e.g. Spain and Italy). For countries with an "extreme high water pressure", the weighting can go up to 56. (See table 31)

Step 2: Valuation

Step 2 a: costs per m³ of water

First, we look at indicators for the marginal impact from 1 m³ of additional use of freshwater, in a region with a medium water scarcity or water stress. In that case, other users will have to look for alternative water sources and/or authorities will oblige other water users to take additional measures for water saving or alternative water sources. We use the additional costs of these alternative water sources and water saving technologies as a proxy for the resource costs of water use.

Table 32 lists a range of different options and technologies that are actually used in countries with moderate to high water stress. We estimate their costs at average 0.15 €/m³, with a range of 0.05 to 0.5 €/m³. This value can also be compared with the central estimate of 0.2 €/m³ used by Alberici (2014) to estimate the external costs of energy use in Europe.

Step 2 b: costs per ecopoint

The above mentioned costs are for a situation in countries with a medium water scarcity and thus weighting factor of 2.25. The costs per ecopoint are thus

Costs per ecopoint = costs per m³ / weighting factor = 0.067

Step 3: Values for different regions and low and high estimates

The variation in water scarcity between regions is accounted for the weighting of ecopoints. So the differences in valuation between regions only account for differences in GDP and sector.

substance	factor	unit
Water, river	0,162	m ³ water eq / m ³
Water, unspecified natural origin, AT	0,012	m ³ water eq / m ³
Water, unspecified natural origin, AU	0,0385	m ³ water eq / m ³
Water, unspecified natural origin, BE	2,84	m ³ water eq / m ³
Water, unspecified natural origin, CA	0,00401	m ³ water eq / m ³
Water, unspecified natural origin, CZ	0,619	m ³ water eq / m ³
Water, unspecified natural origin, DE	1,52	m ³ water eq / m ³
Water, unspecified natural origin, DK	0,736	m ³ water eq / m ³
Water, unspecified natural origin, ES	1,66	m ³ water eq / m ³
Water, unspecified natural origin, extreme water stress	36,8	m ³ water eq / m ³
Water, unspecified natural origin, FI	0,0082	m ³ water eq / m ³
Water, unspecified natural origin, FR	0,619	m ³ water eq / m ³
Water, unspecified natural origin, IT	0,87	m ³ water eq / m ³
Water, unspecified natural origin, low water stress	0,0401	m ³ water eq / m ³
Water, unspecified natural origin, LU	2,84	m ³ water eq / m ³
Water, unspecified natural origin, medium water stress	1,47	m ³ water eq / m ³
Water, unspecified natural origin, moderate water stress	0,368	m ³ water eq / m ³
Water, unspecified natural origin, MX	0,468	m ³ water eq / m ³
Water, unspecified natural origin, NL	0,124	m ³ water eq / m ³
Water, unspecified natural origin, OECD	0,162	m ³ water eq / m ³
Water, unspecified natural origin/kg	0,000162	m ³ water eq / kg
Water, unspecified natural origin/m ³	0,162	m ³ water eq / m ³
Water, unspecified, very high water stress	10,4	m ³ water eq / m ³
Water, well, in ground	0,162	m ³ water eq / m ³

Table 30: Examples of weighting factors for water scarcity for selected countries

Source: Frischknecht, 2006

The values for Flanders/Belgium are adapted accounting for a higher GDP compared to EU 27 (GDP Belgium = 1.19 x GDP EU 27).

The values for the world account for differences in GDP compared to EU 27. In addition, we assume that in general, water use outside the EU is likely to be dominated by agricultural sectors, and that for this sector the costs for water savings and alternative supplies are lower. Therefore we estimate the costs are 50 % lower.

The range of values in the table on costs (table 32) suggests a bandwidth of 9

Low value = central value / $\sqrt{9}$ = central value / 3

High value = central value * $\sqrt{9}$ = central value * 3

Country	Type of measure	Costs In €/m ³	Source
Belgium, Flanders,	Additional costs for drinking water companies to import water from other regions or companies	0.2 (0.05 – 0.35)	Aquaflanders, 2012
Belgium, Flanders,	tax paid by users of freshwater, as a proxy for resource costs	0.04 (0.02-0.06)	
Germany	Incentive paid by water companies to improve water infiltration	0.05	Greiber et al., 2009
France	Incentive paid by private company to improve water infiltration (Bionade)	0.17	VITO
Spain	Costs of water saving measures	0.12-0.21	WWF, cited in Ecologic, 2008
Spain	Costs of water transfer from other basins	0.5 0.3-0.7	
US	Reuse of waste water	0.75 -15	UNEP, 2006
Spain	Building dams to ensure water use	0.06	
France	Making water irrigation systems more efficient (from gravity to pressurized)	0.05	
Spain	Extra fee for extra water use, within a more sophisticated water irrigation system with individual quota and counters	0.06	Cited in Ecologic, 2008
Flanders, Belgium	Additional costs for industrial users that switch from groundwater to 'grey water' (treated waste water)	0.5	
Flanders, Belgium	Additional costs to produce drinking water based on waste water re-use (infiltration of treated waste water)	0.5	
Malta	Costs of producing drinking water based on desalination	1.3	
Europe	Costs of alternative supply measures in EPS valuation	0.003- 0.03	Steen, 2000

Table 31: Costs of different options for water saving and alternative water supply

5.7 Land use occupation

5.7.1 Monetary value

Land use occupation		Value € / kg C deficit		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	1.4E-06	3.4E-07	0.6E-05
	Flanders / Belgium	1.4E-06	3.4E-07	0.6E-05
	Rest of world	1.4E-06	3.4E-07	0.6E-05
Indicators related to biodiversity				
Indicator b1 : loss of ecosystem services for land use urban and industry		Value € / m ² a		
Indicator	Region	Central	Low	High
land use urban, industry, traffic	Western Europe	0.30	0.07	2.35
	Flanders / Belgium	0.59	0.15	2.35
	Rest of world	n.a.	n.a.	n.a.
Indicators b2 and b3 : loss of ecosystem services for land use agriculture and forestry		Value € / m ² a		
Indicator	Region	Central	Low	High
land use agriculture	Western Europe	6.0E-03	1.5E-03	2.4E-02
	Flanders / Belgium	6.0E-03	1.5E-03	2.4E-02
	Rest of world	n.a.	n.a.	n.a.
land use forestry	Western Europe	2.2E-04	5.5E-05	8.8E-04
	Flanders / Belgium	2.2E-04	5.5E-05	8.8E-04
	Rest of world	n.a.	n.a.	n.a.

Table 32: Monetary indicators for land use occupation

n.a. = not available

Source: VITO 2014, update 2017


5.7.2 Methods and data used

The impacts from land occupation and land transformation are very diverse. Impact assessment of land use is still in the first stages of development, capturing the most important impacts (Mila I Canals, 2007; 2008). We distinguish four different impacts.

5.7.2.1 Soil quality and soil organic matter

The first indicator relates to the loss of soil quality. The loss of soil organic matter is a major concern in EU and the rest of the world, both from a perspective of climate change and from loss of soil quality (Smith et al, 2007; EEA, 2012; Schils, 2008, Louwagie, 2009). Loss of soil organic carbon is valued based on the prevention costs, related to measures in agricultural sector to prevent the losses, e.g. zero tillage.

It has been estimated that a carbon tax at the equivalent of 100 \$ per ton CO₂ equivalent would ensure that the most efficient measures will be implemented, and that for cropland and grazing land about half of the technical potential measures will be taken (figure 1). As we value global warming emissions at 100 euro/ton CO₂eq, we also use this cost indicator for the valuation of loss of soil quality. As loss of soil quality is expressed in euro/kg C deficit, we have to correct for the different units (1 kg of CO₂ = 0.27 kg of C).



Figuur 1: Illustration of the mitigation potential for worldwide reduction of GHG measures in agriculture

Source: based on Smith, 2007

The uncertainties related to the valuation of soil quality are inherent to the valuation of carbon as a greenhouse gas. The variation of soil quality impacts depend of measures between countries and different kinds of land uses. We therefore use a bandwidth of 16.

A bandwidth of 16 between our minimum and maximum estimate means that

Low estimate = central estimate / $\sqrt{16}$ = central estimate / 4

High estimate = central estimate * $\sqrt{16}$ = central estimate * 4

5.7.2.2 Biodiversity

Land occupation (urban and industrial)

The impact of land use changes on biodiversity is a very complex issue, and there are no direct tools or data to quantify and monetise these impacts in a systematic way that is relevant for different locations, land uses and species. We therefore follow the approach from TEEB (the economics of ecosystems and biodiversity) (TEEB, 2010). In this worldwide initiative, supported by the EC, land use changes are assessed by accounting for the quantity and value of the ecosystems goods and services from different land uses. They include production services (e.g. agricultural products or wood), regulation services (e.g. carbon sequestration, pollination, etc.) and cultural services (recreation, health, etc.) (see table 34 for the complete list). Whereas production services are mostly valued in private markets, regulation and cultural services are not. Although these services are not a direct indicator for biodiversity, more natural land uses are often associated with both more biodiversity and more regulation and cultural services (Schneiders, 2012). The concept of ecosystem services is used by the Flemish government for the report on the state of nature in Flanders (INBO, 2014)

Land use occupation for urban, industry or transport will result in a loss of unpriced ecosystem services that are associated with open, green spaces. These ecosystem services relate to regulation functions such as removal of pollutants from air by trees and grassland, reduction of noise, global climate regulation (storage of C in biomass), water quality (denitrification). Second, they relate to the cultural services such as recreation, a more pleasant living environment and

health benefits. The loss of production functions (agricultural products and wood) will have been compensated when the land was purchased for industry or urban occupation. The loss of other services has not been compensated.

These services have been identified and valued for open green areas in Flanders. The average for a mixed land uses (agriculture with forests and green areas) is 5900 euro/ha.year. These values represent the situation in Flanders, with on average high levels of pollutants (e.g. air pollution) and a population density three times higher compared to the average for the EU. To estimate a value for Europe, we have corrected these estimates for Flanders, accounting for differences in population densities (see table 34).

Ecosystem service	Flanders (€/ha) (A)	Europe (€/ha) (B)
Production services		
<i>production agricultural products</i>	pm	pm
<i>wood production</i>	pm	pm
Regulation services		
<i>Air quality: capture by plants</i>	1.950	635*
<i>Carbon sequestration in biomass</i>	625	625
<i>Carbon sequestration in soils</i>	946	946
<i>Noise nuisance reduction</i>	0,2	0.1*
<i>Flood prevention</i>	na	na
<i>Water supply</i>	13	13
<i>Nutrient removal</i>	261	85*
<i>Erosion prevention</i>	na	na
<i>Pollination</i>	na	na
Cultural services		
<i>Recreation and tourism</i>	1.400	456*
<i>Quality of living environment</i>	135	44*
<i>Health effects of contact with nature</i>	554	180*
Total (€/ha.a)	5.884	2.984
Total (€/m ² .a)	0,59	0,30

Table 33: Ecosystem services delivered by open green areas in Flanders and Europe

na: not available, pm (pro memory, not to be included to value loss of land use)

* adapted from values from Flanders, accounting for differences in population density.

Source: based on Broekx, 2013

Expressed per ha of land use, these values are similar to the estimates in MMG2012, based on valuation of PDF. The current estimate is more reliable as it avoids building on a single study for valuation of species restoration and the assumption that all species have the same value.

The assessment of ecosystem services includes a wide range of different impacts, and uncertainties for assessment and valuation are high. In addition, for each service, the local situation (soil and habitat type, population density, etc.) may vary significantly. To express both uncertainty and variation, we use a bandwidth of 16.

A bandwidth of 16 between our minimum and maximum estimate means that:

Low estimate = central estimate / $\sqrt{16}$ = central estimate / 4

High estimate = central estimate * $\sqrt{16}$ = central estimate * 4

We do not have comparable data to make reliable estimates for impacts in the rest of the world.

Land occupation (agriculture and forestry)

For agricultural land use we use the prevention costs approach, based on the costs of the measures related to the implementation of target 2 of the EU biodiversity strategy, which is that “By 2020, ecosystems and their services are maintained and enhanced by establishing green infrastructure and restoring at least 15% of degraded ecosystems”. This will require additional measures for land used by agriculture. The costs of these measures have been estimated by Tucker (2013).

From this study, the average cost, expressed per ha of agricultural land (cropland and grassland), is 60 €/ha. We use this figure for the valuation of biodiversity loss on agricultural land.

For forests, this value is estimated at 2.2 €/ha.year.

Uncertainty range:

The authors stress that these numbers are minimum estimates, and the costs per ha may vary a lot depending on country, type of land use and measures required. To express both uncertainty and variation, we use a bandwidth of 16.

A bandwidth of 16 between our minimum and maximum estimate means that

Low estimate = central estimate / $\sqrt{16}$ = central estimate / 4

High estimate = central estimate * $\sqrt{16}$ = central estimate * 4

We do not have comparable data to make reliable estimates for impacts in the rest of the world.

5.8 Land use transformation

5.8.1 Monetary value

Land use transformation		Value € / kg C deficit		
Indicator	Region	Central	Low	high
Full life cycle	Western Europe	1.4E-06	3.4E-07	0.6E-05
	Flanders / Belgium	1.4E-06	3.4E-07	0.6E-05
	Rest of world	1.4E-06	3.4E-07	0.6E-05
Indicators related to biodiversity				
Indicator b1 : transformation from urban land		Value € / m ²		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	n.a.	n.a.	n.a.
	Flanders / Belgium	n.a.	n.a.	n.a.
	Rest of world	n.a.	n.a.	n.a.
Indicator b2 : transformation from agricultural land		Value € / m ²		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	n.a.	n.a.	n.a.
	Flanders / Belgium	n.a.	n.a.	n.a.
	Rest of world	n.a.	n.a.	n.a.
Indicator b3 : transformation from forest, excl. rainforest		Value € / m ²		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	n.a.	n.a.	n.a.
	Flanders / Belgium	n.a.	n.a.	n.a.
	Rest of world	n.a.	n.a.	n.a.
Indicator b4 : transformation from tropical rainforest		Value € / m ²		
Indicator	Region	Central	Low	High
Full life cycle	Western Europe	27*	6.9*	110*
	Flanders / Belgium	27*	6.9*	110*
	Rest of world	27	6.9	110

Table 34: Monetary indicator for land use transformation

n.a. = not available

* Rest of the World monetary indicator are used as default for Western Europe and Flanders

Source: VITO 2014, update 2017

5.8.2 Methods and data used

5.8.2.1 Soil organic matter (C):

The data are the same as for land occupation.

5.8.2.2 Biodiversity: transformation from rainforest (b4)

The transformation of tropical rainforest leads to a loss of valuable areas for biodiversity, with high biodiversity values and very long restoration times. Based on the TEEB study, this loss can be estimated at 8000 €/ha.year or 0.8 €/m².year (Baat, 2008, Chiabai, 2011). To calculate the present value of these annual losses, we have to make assumptions about the discount rate. For the central estimate, we use a discount rate of 3 %, which leads to an indicator of 27.45 €/m².

These numbers are very uncertain, and we use a bandwidth of 16.

Geographically, loss of tropical rainforest is not applicable within Europe or Flanders. However, as within the MMG web tool only the monetary values for Western Europe are taken into account and the other monetary values cannot be chosen, the Rest of the World monetary indicator for transformation from tropical rainforest is used as default for the regions Western Europe and Flanders.

Bijlage 1: List of tables

Table 1: Selected CEN environmental indicators, the corresponding units and LCIA models	10
Table 2: Selected CEN+ environmental indicators, the corresponding units and LCIA models	11
Table 3: Overview of West-European monetary (central, low and high) values for the CEN indicators	11
Table 4: Overview of West-European monetary (central, low and high) values for the CEN+ indicators	12
Table 5: Overview of available methods per impact category and methods used for central, low and high estimations	14
Table 6: Approach to uncertainty in MMG2012 and MMG2014.	18
Table 7: Overview of the uncertainty values for all CEN and CEN+ indicators	19
Table 8: Comparison of indicators for monetary valuation in OVAM MMG and Dutch MMG	20
Table 9: Comparison of indicators for monetary valuation in OVAM MMG and Dutch MMG	21
Table 10: Monetary indicator for depletion of the stratospheric ozone layer	23
Table 11: Monetary indicator for acidification of land and water sources	24
Table 12: Monetary indicator for eutrophication	25
Table 13: Monetary indicator for formation of tropospheric ozone photochemical oxidants	26
Table 14: Monetary indicator for abiotic depletion of non-fossil resources	27
Table 15: Resource depletion costs (RDC) for different substances, based on characterization factor ILCD (method 1)	29
Table 16: Resource depletion costs (RDC) for different substances, based on ratio to market prices (method 2)	30
Table 17: Monetary indicator for abiotic depletion of fossil resources	30
Table 18: Monetary indicator for human toxicity cancer	33
Table 19: Monetary indicator for human toxicity non-cancer	35
Table 20: Monetary indicator for Particulate matter	36
Table 21: Damage costs per kg emitted, for primary particulate matter and particulate matter precursors	38
Table 22: Impacts per ton emitted, for primary particulate matter and particulate matter precursors	39
Table 23: Valuation of health impacts, DALY (disability adjusted life years)	39
Table 24: Impacts from emissions calculated by EcoSense: for year 2015, in EU 27 and Belgium	39
Table 25: Monetary indicator for ionising radiation, human health	40
Table 26: Valuation of DALY for ionising radiation, for Western Europe	41
Table 27: Monetary indicator for ionising radiation on ecosystems	41
Table 28: Monetary indicator for Ecotoxicity, freshwater	42
Table 29: Correction factors used for valuation of Ecotoxicity, freshwater	43
Table 30: Monetary indicator for Water use	43
Table 31: Examples of weighting factors for water scarcity for selected countries	45
Table 32: Costs of different options for water saving and alternative water supply	46
Table 33: Monetary indicators for land use occupation	47
Table 34: Ecosystem services delivered by open green areas in Flanders and Europe	49
Table 35: Monetary indicator for land use transformation	51

Bijlage 2: List of figures

Figuur 1: Illustration of the mitigation potential for worldwide reduction of GHG measures in agriculture

48

Bijlage 3: Bibliography

- Adamowicz, W.L., D. Dupont, A. Krupnick, and J. Zhang. 2011. Valuation of cancer and microbial disease risk reductions in municipal drinking water: an analysis of risk context using multiple valuation methods." *Journal of Environmental Economics and Management*. 61: 213-226
- Anderson, Jason; Kathryn Arblaster; Justin Bartley et al. (2008), *Climate change-induced water stress and its impacts on natural and managed ecosystems*, Study for European Parliament, Ecologic, Berlin, 2008
- Aguilera Roberto F; Roderick G Eggert; Gustavo C C Lagos; John E Tilton (2009), *Depletion and the Future Availability of Petroleum Resources*, *The Energy Journal*; 2009; 30, 1; ABI/INFORM Global, pg. 141
- Aguilera Roberto F; Roderick G Eggert; Gustavo C C Lagos; John E Tilton (2012), *Is Depletion Likely to Create Significant Scarcities of Future Petroleum Resources?*, *Non-Renewable Resource Issues*, *International Year of Planet Earth 2012*, pp 45-82
- Alberici S., Sil Boeve, Pieter van Breevoort, Yvonne Deng, Sonja Förster, Ann Gardiner, Valentijn van Gastel, Katharina Grave, Heleen Groenenberg, David de Jager, Erik Klaassen, Willemijn Pouwels, Matthew Smith, Erika de Visser, Thomas Winkel, Karlien Wouters (2014), *Subsidies and costs of EU energy, An interim report*, Study for the European Commission, Directorate-General for Energy, Ecofys, 2014
- Allacker K., De Troyer F. Geerken T., Spirinckx C., Debacker W., Van Dessel J., Janssen A., Delem L., 2011, *Sustainability, Financial and Quality Evaluation of Dwelling Types – SuFiQuaD*, Final Report, study for the Belgian Science Policy Office (Belspo).
- Allacker K., Souza D.M., Sala S. (2014) *Land use impact assessment in the construction sector: An analysis of LCIA models and case study application*. *The International Journal of Life Cycle Assessment*, *The International Journal of Life Cycle Assessment*, 19(11), pp 1799-1809.
- Ackerman F. and Elizabeth A. Stanton (2012), *Climate Risks and Carbon Prices: Revising the Social Cost of Carbon*. *Economics: The Open-Access, Open-Assessment E-Journal*, Vol. 6, 2012-10. <http://dx.doi.org/10.5018/economics-ejournal.ja.2012-10>
- Anthoff David, Cameron Hepburn, Richard S.J. Tol (2009) , *Equity weighting and the marginal damage costs of climate change*, *Ecological Economics*, Volume 68, Issue 3, 15 January 2009, Pages 836-849, ISSN 0921-8009, <http://dx.doi.org/10.1016/j.ecolecon.2008.06.017>.
- Bickel, P. Friedrich, R., (Eds.), 2001. *Environmental External Costs of Transport*. Springer, Amsterdam.
- Braat, L. and Ten Brink, P. (eds.) with J. Bakkes, K. Bolt, I. Braeuer, B. ten Brink, A. Chiabai, H. Ding, H. Gerdes, M., Jeuken, M. Kettunen, U. Kirchholtes, C. Klok, A. Markandya, P. Nunes, M. van, Oorschot, N. Peralta-Bezerra, M. Rayment, C. Travisi, M. Walpole (2008). *The Cost of Policy Inaction. The case of not meeting the 2010 biodiversity target*. Report of the COPI project, Wageningen and Brussels, May 2008.
- Brin C. en van Grinsven H. (2011), *Costs and benefits of nitrogen in the environment*, in Sutton M.A. (2011), *The European Nitrogen Assessment*, Cambridge University Press, 2011, pp. 513-540

Broekx S., De Nocker L., Liekens I., Poelmans L., Staes J., Van der Biest K., Meire P., Verheyen K. (2013). Raming van de baten geleverd door het Vlaamse Natura 2000-netwerk. Studie uitgevoerd in opdracht van het Agentschap voor Natuur en Bos (ANB/IHD/11/03) door VITO, Universiteit Antwerpen en Universiteit Gent. Brussel. 2013/RMA/R/87

Broekx Steven, Meynaerts Erika, Vercaemst Peter, 2008. Milieukostenmodel Water voor Vlaanderen. Berekeningen voor het stroomgebiedbeheerplan 2009. Studie uitgevoerd in opdracht van het Vlaams Gewest 2009/RMA/R/146.

Broekx S, Smets S, Liekens I, Bulckaen D, Smets S, De Nocker L, Designing a long-term flood risk management plan for the Scheldt estuary using a risk based approach,

Natural Hazards 2011, 57-2, p. 245-266.
<http://www.springerlink.com/content/e43138836415t02n/>

Chiabai, A.; Chiara M. Traversi, Anil Markandya, Helen Ding, Paulo A. L. D. Nunes (2011), Economic Assessment of Forest Ecosystem Services Losses: Cost of Policy Inaction, Environmental and Resource Economics, November 2011, Volume 50, Issue 3, pp 405-445

Cropper M. , James K. Hammitt, Robinson (2011), Valuing Mortality Risk Reductions, Progress and Challenges, RFF discussion paper11-10, Resources for the Future, Washington, 2011

De Bruyn S.M, Korteland M.H, Markowska A.Z., Davidson M.D., de Jong F.L., Bles M., Sevenster M.N., 2010, Handboek Schaduwrijzen – Waardering en weging van emissies en milieueffecten, CE Delft, Delft

Debacker, W., Allacker, K., De Troyer, F., Janssen, A., Delem, L., Peeters, K., De Nocker, L., Spirinckx, C., Van Dessel, J. (2012), Milieugerelateerde Materiaalprestatie van Gebouwelementen, finaal rapport (Environmental Material Performance of Building Elements e Final Report). Public Waste Agency of Flanders (OVAM), available through http://www.ovam.be/sites/default/files/FILE1349102121400ovor121001_MMG_eindrapport.pdf (accessed July 2014).

Debacker W., Allacker K., Spirinckx C., Geerken T., De Troyer F. (2013), Identification of environmental and financial cost efficient heating and ventilation services for a typical residential building in Belgium, in Journal of Cleaner Production, Vol 57, 188-199

Delem L., Van Dessel J., Debacker W., Spirinckx C., Allacker A., De Troyer F., 2011, Bepalingsmethode Milieugerelateerde Materiaalprestatie van gebouwelementen (MMG), voorbereidend rapport voor workshop, in opdracht van de OVAM

De Nocker et al, 2010; Actualisering van de externe milieuschadetekosten (algemeen voor Vlaanderen) met betrekking tot luchtverontreiniging en klimaatverandering, Studie uitgevoerd in opdracht van VMM, MIRA. www.milieurapport.be

De Nocker L., S. Broekx, I. Liekens (2011), Economische waardering van verbetering ecologische toestand oppervlaktewater op basis van onderzoeksresultaten uit Aquamoney, Vito intern rapport 2011/RMA/R/248, 2011, 58 p.

Desaigues, B., Ami, D., Bartczak, A., Braun-Kohlova, M., Chilton, S., Czajkowski, M., Farreras, V., Hunt, A., Hutchinson, M., Jeanrenaud, C., Kaderjak, P., Maca, V., Markiewicz, O., Markowska, A., Metcalf, H., Navrud, S., Nielsen, J. S., Ortiz, R., Pellegrini, S., Rabl, A., Riera, R., Scasny, M., Stoeckel, M. -E., Szanto, R. and Urban, J., 2011. Economic valuation of air pollution mortality: A 9-country contingent valuation survey of value of a life year (VOLY). Ecological Indicators, 11 (3), pp. 902-910

Desaigues B., A. Rabl, D. Ami, Boun My Kene, S. Masson, M-A Salomon and L. Santoni. 2007. "Monetary Value of a Life Expectancy Gain due to Reduced Air Pollution: Lessons from a Contingent Valuation". *Revue d'Economie Politique*, Vol.117 (5), 675-698.

Deutsch F., Veldeman N., Vankerkom J., Peelaerts W., Buekers J., Torfs R., Fierens F., Vanpoucke C., Trimpeneers E., Vancraeynest L., Bossuyt M. (2010), *Zwevend stof en fotochemische luchtverontreiniging. Visionair scenario Milieuverkenning 2030, studie uitgevoerd in opdracht van de Vlaamse Milieumaatschappij, MIRA, MIRA/2010/09, VITO.*

EC, 1996, *ExternE, Externalities of Energy, Volume 2: Methodology*, Luxembourg : European Commission, 1996

EC, 2005, 5, *ExternE - Externalities of Energy – Methodology 2005; Update*. EUR 21951 EN. European Commission, Office for Official Publications of the European Communities, Luxembourg. 2005, ISBN 92-79-00423-9.

EC (2007), *Report from the Commission to the European Parliament and the Council on Addressing the challenge of water scarcity and droughts in the European Union*, SEC(2007) 993

Edenhofer, Ottmar, Brigitte Knopf, Terry Barker, Lavinia Baumstark, Elie Bellevrat, Bertrand Chateau, Patrick Criqui, et al. (2010). *The Economics of Low Stabilization: Model Comparison of Mitigation Strategies and Costs*. *Energy Journal* 31 (Special Issue 1): 11–48. doi:10.5547/ISSN0195-6574-EJ-Vol31-NoSI-2.

EEA, 2009 , *Water resources across Europe — confronting water scarcity and drought*, European environmental Agency, Copenhagen, Report No 2/2009,

EEA, (2011), *Revealing the costs of air pollution from industrial facilities in Europe*, EEA Technical report No 15/2011.

EEA (2012), *The state of soil in Europe*, Copenhagen, 2012

Eftec (2004), *The Health Benefits of Pollution Control: a review of the literature on mortality and morbidity effects*, Paper prepared for UK Department for Environment, Food and Rural Affairs Workshop on Valuation of Health Benefits of Reductions in Air Pollution, London, 2004

Exiopol (2011), *A new accounting framework using externality data and input-output for policy analysis, bottom-up approach*, FEEM, 2011 (www.feem-project.net/exiopol/)

FAO. 2004. *Water charging in irrigated agriculture. An analysis of international experience*. Prepared by G. Cornish, B. Bosworth, C. Perry and J. Burke. *FAO Water Reports*, no. 28.

FEEM, 2008, *CASES project, Cost Assessments for Sustainable Energy Systems*,

<http://www.feem-project.net/cases/>

Flury K. , Niels Jungbluth, Rolf Frischknecht, Ivan Muñoz (2012), *Recommendation for Life Cycle Inventory Analysis for Water Use and Consumption*, Working paper, ESU-services Ltd., Zürich

Frischknecht R., Steiner R., Braunschweig A., Egli N. and Hildesheimer G. (2006) *Swiss Ecological Scarcity Method: The New Version 2006*. In proceedings from: *The Seventh International Conference on EcoBalance*, Nov 14-16, 2006, Tsukuba, Japan, retrieved from: www.esu-services.ch/projects/ubp06/.

Goedkoop M. en Spriensma R. (2000) , The Eco-indicator 99 A damage oriented method for Life Cycle Impact Assessment, Methodology Report, Second Edition, Pré Consultants, 2000, 142 p.

Goedkoop M, Heijungs R., Huijbregts M., De Schrijver A., Struijs J., van Zelm R., 2008, ReCiPe 2008 – A life cycle impact assessment method which comprises harmonised category indicators at the midpoint and the endpoint level – First edition – Report 1: characterisation, in order of Ministerie van Volkshuisvesting, Ruimtelijke Ordening en Milieubeheer.

Gren, I.-M. (2008). Costs and Benefits from Nutrient Reductions to the Baltic Sea. the Swedish Environmental Protection Agency, Stockholm, Sweden.

Hammit, J and Liu, J-T. (2004). Effects of disease type and latency on the value of mortality risk. *Journal of Risk and Uncertainty*. 28.1. 73-95

Hammit JK, Haninger K. (2010). Valuing fatal risks to children and adults: effects of disease, latency, and 1494 risk aversion. *Journal of Risk and Uncertainty* 40:57-83

Hayashi K., Nakagawa A., Itsubo N. and Inaba A. (2006), Expanded Damage Function of Stratospheric Ozone Depletion to Cover Major Endpoints Regarding Life Cycle Impact Assessment, in : *The International Journal of Life Cycle Assessment* , vol. 11, nr. 3 (2006), p. 150–161

Holland, M. , S Pye; G Jones, A Hunt; A Markandya, F Hurley, P Watkiss (2013), The ALPHA Benefit Assessment Model, EC4MACS Task 8 Methodology report, 2013.

Holland, M (2014), Cost-benefit Analysis of Final Policy Scenarios for the EU Clean Air Package, Version 2, Corresponding to IIASA TSAP Report □11, Version 2a, EMRC, October 2014, <http://ec.europa.eu/environment/air/pdf/TSAP%20CBA.pdf>

Hope, Chris A. (2013) ; Critical issues for the calculation of the social cost of CO2: why the estimates from PAGE09 are higher than those from PAGE2002, *J Climatic Change*, V 117, N 3, Springer, Netherlands, 2013

Huijbregts MAJ, LJA Rombouts, AMJ Ragas (2005), Human-toxicological effect and damage factors of carcinogenic and noncarcinogenic chemicals for life cycle impact assessment - Integrated Environmental Assessment and ..., 2005

Huijbregts, U Thissen, T Jager, D Van de Meent, AMJ Ragas (2000), Priority assessment of toxic substances in life cycle assessment. Part II: assessing parameter uncertainty and human variability in the calculation of toxicity potentials, *Chemosphere*, vol 41, Nr. 4, 575-588

Humbert, S, Marshall, J.D, Shaked, S, Spadaro, J.V., Nishioka, Y., Preiss, P., McKone, T.E., Horvath, A. and Jollet, O. (2011), Intake Fraction for Particulate Matter: Recommendations for Life Cycle Impact Assessment. *Environmental Science & Technology*. 45. 4808-4816.

International Energy Agency (2010). *Energy Technology Perspectives 2010: Scenarios & Strategies to 2050*. Paris. Available at <http://www.iea.org/techno/etp/index.asp>.

INBO (2014), *Ecosysteemdienstenrapporten Vlaanderen* verschenen, <http://www.nara.be/>

JRC 2010, European Commission - Joint Research Centre (JRC) - Institute for Environment and Sustainability (IES), *International Reference Life Cycle Data System (ILCD) Handbook - General guide for Life Cycle Assessment - Detailed guidance*. First edition March 2010

Kounina Anna, Manuele Margni, Jean-Baptiste Bayart, Anne-Marie Boulay, Markus Berger, Cecile Bulle, Rolf Frischknecht, Annette Koehler, Llorenç Milà i Canals, Masaharu Motoshita, Montserrat Núñez, Gregory Peters, Stephan Pfister, Brad Ridoutt, Rosalie van Zelm, Francesca Veronesi, Sebastien Humbert (2013), Review of methods addressing freshwater use in life cycle inventory and impact assessment, The International Journal of Life Cycle Assessment, March 2013, Volume 18, Issue 3, pp 707-721

Kuik, O., Brandera, L., Nikitina, N., Navrud, S., Magnussen, K., El Hadji, F., 2008. A Database of Studies on Energy-related External Costs due to Land Use Changes, Acidification and Eutrophication, Visual Intrusion and Climate Change. Delivery of FP6 Cases Project

Le Goffe, P. (1995): "The Benefits of Improvements in Coastal Water Quality: A Continental Approach", Journal of Environmental Management 45, (4): 305-317.

Liekens I., De Nocker L., 2008. Rekenraamwerk voor de economische baten van een betere waterkwaliteit, VITO in opdracht van VMM, september 2008. <http://www.milieurapport.be> onderzoeksrapporten 2008

Liekens I., Schaafsma M., Staes J., Brouwer Roy, De Nocker L. en Meire P., 2009. Uitvoeren van een economische waarderingsstudie van natuurlandschappen voor gebruik in MKBA van infrastructuurprojecten. Rapport in opdracht van LNE, afdeling milieu-, natuur- en energiebeleid. VITO, 2009/RMA/R/308.

Luengo-Fernandez Ramon, Jose Leal, Alastair Gray and Richard Sullivan (2013): Economic burden of cancer across the European Union: a population-based cost analysis. - The Lancet Oncology, 2013 Nov;14(12):1165-74.

Louwagie, G; Hubertus Gay, Alison Burrell (2009), Final report on the project 'Sustainable Agriculture and Soil Conservation, JRC 51775

Lloyd Shannon M., Robert Ries (2008), Characterizing, Propagating, and Analyzing Uncertainty in Life-Cycle Assessment A Survey of Quantitative Approaches, Journal of Industrial Ecology, 2008, 11, 1, 161-179

LNE (2007), milieubeleidskosten – begrippen en berekeningsmethoden, Vlaamse Overheid, LNE, 2007, www.milieueconomie.lne.be, p. 27 e.v.

LNE, 2008 Milieubaten of milieuschadetekosten-waarderingsstudies in Vlaanderen, Vlaamse overheid, Departement LNE, 2008, www.milieueconomie.lne.be

Maibach M., C. Schreyer, D. Sutter ; H.P. van Essen, B.H. Boon, R. Smokers, A. Schroten ; C. Doll ; B. Pawlowska, M. Bak (2008), Handbook on estimation of external costs in the transport sector. Internalisation Measures and Policies for All external Costs of Transport (IMPACT), Version 1.1, Delft, CE Delft, 2008

Mariotto AB, Yabroff KR, Shao Y, Feuer EJ, and Brown ML. Projections of the Cost of Cancer Care in the United States: 2010-2020. JNCI, Jan 19, 2011, Vol. 103, No. 2

Markandya, A., Nunes, P.A.L.D., Brauer, I., ten Brink, P. Kuik, O. and M. Rayment (2008) "Review On The Economics Of Biodiversity Loss – Economic Analysis and Synthesis", Final report for the European Commission, Venice, Italy. 140 pp.

McKinsey & Company (2009). Pathways to a Low-Carbon Economy: Version 2 of the Global Greenhouse Gas Abatement Cost Curve. http://www.mckinsey.com/clientservice/ccsi/pathways_low_carbon_economy.asp.

MEA (2005). Ecosystems and Human Well-being: Biodiversity Synthesis, World Resources Institute, Ashington, D.C.

Milà i Canals L, Bauer C, Depestele J, Dubreuil A, Freiermuth Knuchel R, Gaillard G, Michelsen O, Müller-Wenk R, Rydgren B (2007) Key elements in a framework for land use impact assessment in LCA. *Int J Life Cycle Assess* 12(1):5–15

Milà i Canals L, Romanyà J, Cowell SJ (2006): Method for assessing the use of 'fertile land' in Life Cycle Assessment (LCA). *Journal of Cleaner Production* 15 (2007) 1426-1440

MIRA (2007) Milieu- en natuurrapport Vlaanderen, Achtergronddocument 2007 Vermesting. Overloop S., Bossuyt M., Ducheyne S., Dumortier M., Eppinger R., Van Gijsegem D., Van Hoof K.,

NDCF, (2007), The hidden costs of oil, an update, National Defence Council Foundation, 2007

NIH (2011), National Cancer Institute. "Cancer costs projected to reach at least \$158 billion in 2020."

NIH (2013), National Institutes of Health. Center to Reduce Cancer Health Disparities (2013) . Economic Costs of Cancer Health Disparities. <http://crchd.cancer.gov/attachments/NCIeconomiccosts.pdf>. Accessed oct. 2013.

Ott et al, 2006 , Ott, Walter, Baur, Martin and Kaufmann, Yvonne, e c o n c e p t AG, Frischknecht, Rolf and Steiner, Roland, ESU-services, 2006. , NEEDS, project, FP6, RS1b_D4.2, "Assessment of Biodiversity Losses".

Preiss P., Anna Kounin, Sandra Torras, Joachim Roos, Rita Van Dingenen, Sebastien Humbert, Olivier Jolliet, Frank Dentener, Rainer Friedrich (2013), Recommended assessment framework, method and characterization factors for human health impacts of fine particulate matter formation; Development and application of environmental Life Cycle Impact assessment Methods for improved sustainability Characterisation of Technologies, LC Impact project, Stuttgart, 2013

Preiss, Ph, Volker Klotz (2008a), revised Description of updated and extended tools for the detailed site-dependent assessment of External costs EcoSenseWeb, Report NEEDS, IER, 2008;

Pretty ,J. N.,Mason,C. F.,Nedwell,D. B. et al. (2003).Environmental costs of freshwater eutrophication in England and Wales. *Environmental Science and Technology*, 37,201 –208.

Rosenbaum R.K., Bachmann T.M., Swirsky Gold L., Huijbregts M.A.J., Jolliet O., Juraske R., Koehler A., Larsen H., MacLeod M., Margni M., McKone T.E., Payet J., Schuhmacher M., van de Meent D., Hauschild M. (2008), USEtox – the UNEP SETAC toxicity model: recommended characterisation factors for human toxicity and freshwater ecotoxicity in life cycle impact assessment, in *International Journal of Life Cycle Assessment*, Volume 13, Issue 7, pp 532-546.

Ruijgrok, Dr.ir. E.C.M. , Dr.ir. R. Brouwer prof. dr. H. Verbruggen, 2004. Waardering van Natuur, Water en Bodem in Maatschappelijke Kosten-batenanalyses. Aanvulling op de Leidraad OEI. In opdracht van het ministerie van Landbouw, Natuur en Voedselkwaliteit, in samenwerking met de ministeries van Verkeer en Waterstaat, Economische Zaken, Financiën en VROM.RWS 2004

Ricardo-AEA (2014), Update of the Handbook on External Costs of Transport, Final Report, commissioned by European Commission – DG Mobility and Transport, 8 January 2014, available through <http://ec.europa.eu/transport/themes/sustainable/studies/doc/2014-handbook-external-costs-transport.pdf>

SBK (2011), Bepalingsmethode Milieuprestatie Gebouwen en GWW-werken, Berekeningswijze voor het bepalen van de milieuprestatie van gebouwen en GWW-werken gedurende hun gehele levensduur, gebaseerd op de levenscyclusanalyse methode, via <https://www.milieudatabase.nl/index.php?id=basisinformatie>

Schneiders A., Toon Van Daele, Wouter Van Landuyt, Wouter Van Reeth, Biodiversity and ecosystem services: Complementary approaches for ecosystem management?, *Ecological Indicators*, Volume 21, October 2012, Pages 123-133, ISSN 1470-160X, <http://dx.doi.org/10.1016/>

René Schils, Peter Kuikman, Jari Liski, Marcel van Oijen, Pete Smith, Jim Webb, Jukka Alm, Zoltan Somogyi, Jan van den Akker, Mike Billett, Bridget Emmett, Chris Evans, Marcus Lindner, Taru Palosuo, Patricia Bellamy, Jukka Alm, Robert Jandl and Ronald Hiederer (2008), Review of existing information on the interrelations between soil and climate change, climsoil project, European commission, Technical Report - 2008 - 048

Smith, P., D. Martino, Z. Cai, D. Gwary, H. Janzen, P. Kumar, B. McCarl, S. Ogle, F. O'Mara, C. Rice, B. Scholes, O. Sirotenko, (2007): Agriculture. In *Climate Change 2007: Mitigation. Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [B. Metz, O.R. Davidson, P.R. Bosch, R. Dave, L.A. Meyer (eds)], Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA

Söderqvist, T. and Hasselström, L. (2008). *the Economic Value of Ecosystem Services Provided by the Baltic Sea and Skagerrak*.

<http://www.naturvardsverket.se/Documents/publikationer/978-91-620-5874-6.pdf>

Stern, N., Peters, S., Bakhshi, V., Bowen, A., Cameron, C., Catovsky, S., Crane, D., Cruickshank, S., Dietz, S., Edmondson, N., Garbett, S., Hamid, L., Hoffman, G., Ingram, D., Jones, B., Patmore, N., Radcliffe, H., Sathiyarajah, R., Stock, M., Taylor, C., Vernon, T., Wanjie, H. & Zenghelis, D. (2006), *Stern review on the economics of climate change*, Cambridge University Press, Cambridge.

TNO (2004), Toxiciteit heeft z'n prijs Schaduw prijzen voor (eco-)toxiciteit en uitputting van abiotische grondstoffen binnen DuboCalc, in order of Rijkwaterstaat (Dutch Ministry of Infrastructure and Environment), 8 maart 2004.

Tol R.S.J., 2009, The Economic Effects of Climate Change, *Journal of Economic Perspectives*—Volume 23, Number 2—Spring 2009—Pages 29–51

Tol, R.S.J. (2011), The Social Cost of Carbon, *Annual Review of Resource Economics*, 3, 419-443.

Tol, R.S.J. (2012), 'The Uncertainty about the Total Economic Impact of Climate Change', *Environmental and Resource Economics*, 53 (1), 97-116.

Torfs R., Hurley F., Miller B., Rabl A. (2007), A set of concentration-response functions. NEEDS, New Energy Externalities Developments for Sustainability, Deliverable 3.7 – RS1b/WP3, 162 pp., Vito, 2007.

Yabroff, K.R., Warren, J.L., Brown, M.L. (2007). Costs of cancer care in the USA: a descriptive review. *Nat Clin Pract Oncol.*, 4(11), 643-56

Van Houtven G, Sullivan MB, Dockins C. 2008. Cancer premiums and latency effects: a risk tradeoff 1657 approach for valuing reductions in fatal cancer risks. *Journal of Risk and Uncertainty* 36(2):179-199

van Vuuren, Detlef P., Elke Stehfest, Michel G. J. Elzen, Tom Kram, Jasper Vliet, Sebastiaan Deetman, Morna Isaac, et al. (2011). RCP2.6: Exploring the Possibility to Keep Global Mean Temperature Increase below 2°C. *Climatic Change* 109 (1–2) (August 5): 95–116.
doi:10.1007/s10584-011-0152-3.

Van Zelm R., Huijbregts M.A.J., Harbers J.V., Wintersen A., Posthuma L., Struijs J., Van de Meent D. (2007) Uncertainty in msPAF-based ecotoxicological effect factors for freshwater ecosystems in life cycle impact assessment. *Integrated Environmental Assessment and Management* 3(2): 203-210; 3(4) e6-e37.

Van Zelm R, Mark A.J. Huijbregts, Hans A. van Jaarsveld, Gert Jan Reinds, Jaap Struijs & Dik van de Meent (2007). Time horizon dependent characterization factors for acidification in life-cycle assessment based on forest plant species occurrence in Europe. *Environmental Science & Technology* 41 (2007), 922-927

Van Zelm R, Huijbregts MAJ, Posthuma L, Wintersen A, Van de Meent D. 2009. Pesticide Ecotoxicological Effect Factors and Their Uncertainties for Freshwater Ecosystems. *Int. J. LCA* 14: 43-51.

Vogels N., Vanden Auweele W., Wustenberghs H., D'hooghe J., Vlaamse Milieumaatschappij, www.milieuraapport.be

Waldhoff S., David Anthoff, Steven Rose, and Richard S. J. Tol (2014), *The Marginal Damage Costs of Different Greenhouse Gases: An Application of FUND*, October 01, 2014

Watkiss, P., Anthoff, D., Downing, T., Hepburn, C., Hope, C., Hunt, A., Tol, R. (2005): *The Social Cost of Carbon (SCC) Review – Methodological Approaches for Using SCC Estimates in Policy Assessment*. Final Report. AEA Technology Environment, Harwell, November 2005

Watkiss, P and Downing, T (2008). *The Social Cost of Carbon: Valuation Estimates and Their Use in UK Policy*. *Integrated Assessment Journal*. Vol. 8, Iss. 1 (2008), Pp. 85–105

Watkiss, P., Anthoff, D., Downing, T., Hepburn, C., Hope, C., Hunt, A. & Tol, R. S. J. (2006), *The social costs of carbon (SCC) review—methodological approaches for using SCC estimates in policy assessment*, Final report to DEFRA.

UK DECC (2011), . A brief guide to the carbon valuation methodology for UK policy appraisal. Department of Energy and Climate Change. (October 2011)
<http://www.decc.gov.uk/assets/decc/11/cutting-emissions/carbon-valuation/3136-guide-carbon-valuation-methodology.pdf>.

UK, DECC (2014), *Valuation of energy use and greenhouse gas (GHG) emissions, Supplementary guidance to the HM Treasury Green Book on Appraisal and Evaluation in Central Government*, Londen, 2014

US DOE (2010), Interagency Working Group on Social Cost of Carbon (2010). *Social Cost of Carbon for Regulatory Impact Analysis Under Executive Order 12866*. Washington, DC: U.S. Department of Energy.
http://www1.eere.energy.gov/buildings/appliance_standards/commercial/pdfs/smallmotors_tsd/em_finalrule_appendix15a.pdf

UK, DECC (2014), *Valuation of energy use and greenhouse gas (GHG) emissions, Supplementary guidance to the HM Treasury Green Book on Appraisal and Evaluation in Central Government*, Londen, 2014

US DOE (2010), Interagency Working Group on Social Cost of Carbon (2010). *Social Cost of Carbon for Regulatory Impact Analysis Under Executive Order 12866*. Washington, DC: U.S.

Department of Energy.

http://www1.eere.energy.gov/buildings/appliance_standards/commercial/pdfs/smallmotors_tsd/em_finalrule_appendix15a.pdf

Waldhoff S., David Anthoff, Steven Rose, and Richard S. J. Tol (2014), The Marginal Damage Costs of Different Greenhouse Gases: An Application of FUND , October 01, 2014

Watkiss, P and Downing, T (2008). The Social Cost of Carbon: Valuation Estimates and Their Use in UK Policy. *Integrated Assessment Journal*. Vol. 8, Iss. 1 (2008), Pp. 85–105

Watkiss, P., Anthoff, D., Downing, T., Hepburn, C., Hope, C., Hunt, A. & Tol, R. S. J. (2006), The social costs of carbon (SCC) review—methodological approaches for using SCC estimates in policy assessment, Final report to DEFRA.

Webster, A., Donald Mackay (2003), Defining Uncertainty and Variability in Environmental Fate Models, CEMC Report No. 200301, Canadian Environmental Modelling Centre, Ontario, 2003

World Health Organization (2012). Measurement of healthy life expectancy and wellbeing: report of a technical meeting, Geneva 10-11 December 2012. Geneva: World Health Organization.